

138

March 21st, 1925.

Vol. XLV., Art. 5.

東京帝國大學
理學部紀要

1925a

第四拾五冊 第五編

(17)

JOURNAL
OF THE
COLLEGE OF SCIENCE,
IMPERIAL UNIVERSITY OF TOKYO.

22112
22364

MATAJIRO YOKOYAMA,

Molluscan Remains from the Uppermost Part of the
Jô-Ban Coal-Field.



四

TOKYO.

PUBLISHED BY THE UNIVERSITY.

TAISHO XIV.

27088
1
27091

Publishing Committee.

Prof. **S. Goto**, *Rigakuhakushi*, Director of the College (*ex officio*).

Prof. **K. Matsubara**, *Rigakuhakushi*.

Prof. **K. Fujii**, *Rigakuhakushi*.

Prof. **N. Yamasaki**, *Rigakuhakushi*.

Prof. **N. Yatsu**, *Ph. D.*, *Rigakuhakushi*.



All communications relating to this Journal should be addressed to the
Director of the College of Science.

Molluscan Remains from the Uppermost Part of the Jō-Ban Coal-Field.

By

Matajiro YOKOYAMA, Rigakuhakushi.

With 6 Plates.

In my paper recently published on the fossils of the lowest part of the Jō-Ban coal-field,¹⁾ I enumerated the several beds of which this coal-field is composed. The present paper treats of the Molluscan remains found in its uppermost beds, the so-called Shirado-Beds, which seem to be most fossiliferous, when compared with the others.

These remains are partly those collected by Dr. Tokunaga in his recent researches in the said coal-field, and partly those kept in the Museum of the Geological Institute of the Imperial University of Tokyo. Among the latter, there are many which have been collected by Dr. T. Kochibe as early as 1879, when he was studying the geology of the region corresponding to the southern part of what we now call the Jō-Ban Coal-field, and which he mentions in his "Geology of Northern Hitachi" published in 1882.²⁾

The number of fossil localities amounts to more than forty, but the greater part of the fossils have been obtained at four places, viz.: Sukegawa, Tempizan, Yunami and Izura, the first having yielded 53 species, the second 23, the third 19, and the fourth 17. The remaining localities, taken all together, have given 42 species, although each one taken separately has yielded only a few, often only a single species. It is much to be regretted that where the fossils are most numerous they are mostly found as casts, thus making their determination not only difficult, but in many cases impossible.

The number of species found in all the localities taken together amounts to 96, as shown in the following table:

1) Molluscan Remains from the Lowest Part of the Jō-Ban Coal-Field. *Jour. Sci. Coll.*, Vol. XLV, Art. 3, 1924.

2) Written in Japanese, the title in Japanese being "Jōhoku Chitsu Hen." *Mem. Tokio Daigaku*, No. 4.

	Living.					
	Sukagawa	Tempizan	Izura	Yamami	Other localities	Musa-shino
I. Mollusca.						
1. Terebra sp.						
2. Conus sp.						
3. Pleurotoma mediocariata Yok.				+	+	+
4. Pleurotoma sp.		+				
5. Drillia principalis Pils.		+			+	Northern, Central and Western Japan.
6. Bela rugulata Tros. var. schneideri Harm.					+	(Fossil in English Crag).
7. Olivella fortunei Ad.	?				+	Central Japan.
8. Voluta megaspira Sow.	+	+		+	+	North., Centr., West. Japan.
9. Chrysodomus phoeniceus Dall.	+	+	+	+	+	Japan? British Columbia.
10. Chrysodomus pericochlion Schr.	+					Northern Japan.
11. Buccinum leucostoma Lke.	+			+	+	Central Japan.
12. Volutharpa perryi Jay.	+				+	Northern and Central Japan.
13. Priene oregonensis Redf.	+				+	Northern Japan.
14. Galeodea (Sconsia) japonica Yok.		+	+			(Pliocene of Izumo).
15. Cassis japonica Rve.	+					Central and Western Japan.
16. Dolium luteostomum Küst.	+				+	North., Centr., West. Japan.
17. Cerithium baculum Yok.					+	(Pliocene of Kii).
18. Cerithium sp.				+	+	
19. Vermetus sp.	+					
20. Turritella nipponica Yok.	+	+	+	+	+	
21. Solarium sp.	+					
22. Calyptraea mammilaris Brod.		+		+	+	West coast of America.
23. Crepidula sp.	+					
24. Natica janthostoma Desh.	+	+	+	+	+	Northern and Central Japan.
25. Polinices (Neverita) ampla Phil.	+				?	North., Centr., West. Japan.
26. Sigaretus (Eunaticina) papilla Gmel.			+			Central and Western Japan.
27. Turbo (Batillus) cornutus Gmel.	+					North., Centr., South. Japan.
28. Bathyembix argenteonitens Lke.	+					Central and Western Japan.
29. Stomatella japonica Ad.	+				+	Western Japan.
30. Haliotis sp.	+					

	Sukegawa	Tenjizan	Enra	Yunami	Other localities	Moss-shimo	Living.
31. <i>Acmaea</i> sp.		+					
32. <i>Helcioniscus pallidus</i> Gld.	+			+		+	Northern and Central Japan.
33. <i>Dentalium weinkauffii</i> Dkr.	+						Central Japan.
34. <i>Dentalium complexum</i> Dall.		+			+	+	Central Japan. Hawai.
35. <i>Panope generosa</i> Gld.				+	+	+	Northern Japan. West. America.
36. <i>Mya arenaria</i> L. var. <i>japonica</i> Jay.				+			North., Centr., West. Japan.
37. <i>Mactra sulcataaria</i> Desh.	+	+				+	" " " "
38. <i>Mactra veneriformis</i> Desh.				+		+	" " " "
39. <i>Mactra spectabilis</i> Lke.			+		+		Central and Western Japan.
40. <i>Spisula grayana</i> Schr.	+	+			+	+	Northern Japan.
41. <i>Solen grandis</i> Dkr.						+	Western Japan.
42. <i>Solen gouldii</i> Conr.				+			Northern and Central Japan.
43. <i>Cultellus izumoensis</i> Yok.					+		(Pliocene of Izumo).
44. <i>Siliqua pulchella</i> Dkr.					+	+	Central and Western Japan.
45. <i>Soletellina violacea</i> Lam.					+	+	North., Centr., West. Japan.
46. <i>Tellina besshoensis</i> Yok.					+		(Miocene of Iwaki).
47. <i>Tellina optiva</i> Yok.					+		(Pliocene of Izumo).
48. <i>Tellina izurensis</i> Yok.		+					
49. <i>Macoma praetexta</i> Mart.	+				+	+	Central and Western Japan.
50. <i>Macoma dissimilis</i> Mart.	+	?	?		+	+	Central Japan.
51. <i>Dosinia troscheli</i> Lke.	+	+			+	+	Central and Western Japan.
52. <i>Cyclina chinensis</i> Chem.					+	+	North., Centr., West. Japan.
53. <i>Meretrix iizukai</i> Yok.	+		+	+	+		
54. <i>Clementia speciosa</i> Yok.					+		(Pliocene of Dainichi).
55. <i>Venus (Mercenaria) stimpsoni</i> Gld.				+		+	North., Cent., West. Japan.
56. <i>Venus jedoensis</i> Lke.	+					+	Central and Western Japan.
57. <i>Venus toreuma</i> Gld.	+	+					" " " "
58. <i>Tapes undulatus</i> Born.	+						" " " "
59. <i>Cardium californiense</i> Desh.		+				+	North., Centr., West. Japan.
60. <i>Cardium muticum</i> Rve.					+	+	" " " "

	Sukagawa	Tempzan	Izuna	Yunami	Other localities	Masa-shiu	Living.
61. <i>Cardium modestum</i> Ad. et Rve.	+					+	Central Japan. (Miocene and Pliocene).
62. <i>Cardium shinjiense</i> Yok.	+	+		+	+		Central and Western Japan.
63. <i>Diplodonta semiaspera</i> Phil.	+		+		+	+	Central Japan.
64. <i>Lucina (Phacoides) borealis</i> L.	+		+	+	+	+	Alaska. North. Pacific. Puget Sound.
65. <i>Thyasira bisecta</i> Conr.	+				+		Central and Western Japan.
66. <i>Cardita cumingiana</i> Dkr.	+					+	Northern Japan.
67. <i>Venericardia ferruginea</i> Ad.				+	+	+	Atlantic. (Fossil in Saishu).
68. <i>Thracia pubescens</i> Pult.					+		North.—South. Japan.
69. <i>Mytilus grayanus</i> Dkr.		+					Central-Southern Japan.
70. <i>Mytilus</i> sp.		+					North., Central Japan, Philip-pines.
71. <i>Modiola</i> sp.	+						Central Japan. Patagonia.
72. <i>Lima lima</i> L.	+						Northern, Central, Western Japan.
73. <i>Lima angulata</i> Low.	+					+	Central Japan. Central Japan.
74. <i>Lima vulgatula</i> Yok.	+					+	Central Japan.
75. <i>Lima goliath</i> Sm.	+	+			+	+	Northern Japan. Sea of Okhotsk.
76. <i>Pecten laetus</i> Gld.	+	+				+	Northern Japan. Sea of Okhotsk.
77. <i>Pecten vesiculosus</i> Dkr.	+					+	Centr., West., South. Japan.
78. <i>Pecten swiftii</i> Bern.	+					+	(Pliocene of Kii).
79. <i>Pecten yessoensis</i> Jay.		+	+		+	+	Central Japan.
80. <i>Pecten kimurai</i> Yok.	+		+	+	+	+	Northern and Central Japan.
81. <i>Ostrea gigas</i> Thunb.		+				+	Centr., West., South. Japan.
82. <i>Pinna japonica</i> Hanl.					+	+	North., Centr., West. Japan.
83. <i>Arca kobeltiana</i> Pils.	+	+				+	Central and Western Japan.
84. <i>Arca subcrenata</i> Lke.	+				+	+	Northern and Central Japan.
85. <i>Arca setoensis</i> Yok.					+	+	Central Japan.
86. <i>Pectunculus vestitus</i> Dkr.		+				+	Northern and Central Japan.
87. <i>Parallelodon obliquatus</i> Yok.	+	+				+	Central Japan.
88. <i>Limopsis azumana</i> Yok.	+					+	China Sea.
89. <i>Leda confusa</i> Hanl.			?		+	+	Central Japan.
90. <i>Leda</i> sp.	+					+	

	Sukagawa	Tempizan	Izura	Yunami	Other localities	Musa-shino	Living.
91. <i>Yoldia lisichkei</i> Smith.	+			+			Central Japan.
92. <i>Nucula insignis</i> Ad.	+					+	Northern Japan.
93. <i>Nucula mirabilis</i> Ad. et Rve.	+					+	Central and Western Japan.
94. <i>Solemya tokunagai</i> Yok.					+		
II. Brachiopoda.							
95. <i>Terebratulina caput-serpentis</i> L.	†					+	Central Japan. North. Pacific. Atlantic.
96. <i>Pereudesia grayi</i> Dav. var. <i>transversa</i> Dav.		+					Northern, Central, Western Japan. California.

Of the 96 species above enumerated, 12 are not well determined, and of the remaining 84, 17 are not yet known as living. They are the following :

1. *Pleurotomma mediocarinata* Yok.
2. *Bela rugulata* Tros., var. *schneideri* Harm.
3. *Galeodea (Scosnia) japonica* Yok.
4. *Cerithium baculum* Yok.
5. *Turritella nipponica* Yok.
6. *Cultellus izumoensis* Yok.
7. *Tellina besshoensis* Yok.
8. *Tellina optiva* Yok.
9. *Tellina izurensis* Yok.
10. *Meretrix iizukai* Yok.
11. *Clementia speciosa* Yok.
12. *Cardium shinjiense* Yok.
13. *Lima vulgaris* Yok.
14. *Pecten kimurai* Yok.
15. *Limopsis azumana* Yok.
16. *Arca setoensis* Yok.
17. *Solemya tokunagai* Yok.

Of these 17, which make about 20% of the whole fauna, 13 have already been found fossil in the same or other parts of Japan; namely, 5 in the Musashino Formation of the neighbourhood of Tokyo, 7 in the Pliocene of Totomi, Kii and Izumo, and 1 in the lowest part (Miocene) of the same coal-field, while 4 are entirely new forms.

Besides the above 17, there are 4 or at least 3 which are not yet known from Japanese waters. They are *Calyptrea mammilaris* Brod. of the west coast of America, *Thyasira bisecta* Conr. of the eastern half of the

Northern Pacific and *Thracia pubescens* Pult. of the Atlantic. *Chrysodonmus phoeniceus* Dall which has been described from the northern part of the west coast of America may possibly be found also in the Japanese seas, as I found a specimen in our Geological Institute with no locality given, but which does not seem to be foreign.

The whole fauna, when compared with the already known ones in Japan, shows closest relationship to that of the *Musashino Formation*. Out of the 84 species which constitute it, 61 or 72.6% occur in this latter formation. Consequently, the Shirado-Beds from which these fossils have been obtained, if they are Tertiary at all, must be looked upon as *Upper Pliocene*, and not older.

Very interesting is the more northern aspect of the fossil fauna, when compared with the recent one of the neighbouring seas. The 67 living species grouped together according to their present habitat are as follows:

1.	Species now living only near the fossil localities (Central Japan) or in about the same latitudes (Western Japan). ... 28	Percentage of the whole fauna. 41.7
2.	Species now living in Central or Western Japan as well as north of it. 22	Percentage of the whole fauna. 32.8
3.	Species now living in Central or Western Japan as well as south of it (Southern Japan and further south). 2	Percentage of the whole fauna. 3.0
4.	Species now living in whole Japan (Northern—Southern Japan). 3	Percentage of the whole fauna. 4.5
5.	Species now living only in Northern Japan... 8	Percentage of the whole fauna. 12.0
6.	Species now living only outside of Japan. 4.0	Percentage of the whole fauna. 6.0

As might be expected from the very young age of the fossil fauna, out of 67 living forms, 55 or 82% are those now living in Central and Western Japan, that is to say, in the seas near the fossil localities or further to the west, but in about the same latitudes. However, if we divide these 55 into four groups (no. 1-no. 4) as above done, the number of species also living in the north far exceeds those also living in the south, being more than ten times

As in my three previous papers, Northern Japan is north of the 38th parallel, north latitude, while Central Japan is the part lying south of it and east of 138° east longitude. Western Japan is the part west of the same longitude, including, Chugoku, Shikoku and Kyushu. Southern Japan includes all the islands lying south of Kyushu.

as many. Besides, what I have to emphasize is the occurrence of 8 species which now live only in Northern Japan. They are the following:

1. *Chrysodomus pericochlion* Schr.
2. *Priene oregonensis* Redf.
3. *Panope generosa* Gld.
4. *Spisula grayana* Schr.
5. *Venericardia ferruginea* Ad.
6. *Pecten swiftii* Bern.
7. *Pecten yessoensis* Lay.
8. *Nucula insignis* Ad.

To these 4 others are to be added which have the same value as these north-Japanese forms. They are the two deep-sea ones *Lima goliath* Sm. and *Yoldia lischkei* Sm. (above included in no. 1) and two foreign ones *Chrysodomus phoeniceus* Dall and *Thyasira bisecta* Conr. These altogether make 14, which is about 20% of the living species.

From this it becomes quite evident that the fossil fauna represents one which is more northern or, as naturalists usually say, more boreal than the recent. And this is quite in accordance with the results¹⁾ obtained by studying the Musashino fossils. Therefore, so far as the molluscan remains are concerned, it now becomes imperative to assume a more or less cooler state of the sea-water of Central Japan during the later Pliocene epoch, compared with the present.²⁾

1) Yokoyama. Fossils from the Miura Peninsula and its Immediate North, p. 24. Fossils from the Upper Musashino of Kazusa and Shimosa, p. 21.

2) Among our savants, there is one who, for several years, has been trying to negate, or at least to belittle, the conclusion at which I have arrived on the faunistic character of the Musashino fossils, by purposely studying a group of Protozoa called Foraminifera found in the same formation. Whatever may be the result which he draws from these minute creatures, the one obtained by me from the Musashino Mollusca remains unshaken.

By the way, it seems to me that this endeavor has arisen from a misunderstanding, as if I had taken the Musashino fauna for "arctic," because this savant says that the Foraminifera taken altogether are not at all "arctic." It is true that I have pointed out the intermingling of a few arctic species among the fauna, but the fauna itself I never called "arctic" on that account. I simply said that it bears a "more boreal character, when compared with the recent" ("Fossils from the Miura Peninsula, p. 22). By the term arctic," as I understand it, is meant around the "North Pole," while "boreal" means simply "northern" and no more.

Furthermore, I must say that his statement that I consider the Lower Musashino as *Lower Pliocene* is not quite correct. I said in the work above quoted, p. 21, that the fauna is to be brought somewhere near that of the Red Crag of Newbourne (Newbourian). If this Newbourian is according to his own opinion *Lower Pliocene*, I have nothing to say. But if he accepts the classification of the Crag generally adopted in Europe, he should call it *Upper Pliocene*, or at lowest *Middle Pliocene*.

I must not omit here to mention the uncommonly large growth of the two fossil species of *Lucina (Phacoides) borealis* and *Venericardia ferruginea*, in comparison to their recent representatives. The large specimens of the former, I have already noticed in studying the Musashino fossils, which I then took as belonging to a distinct species. But by examining the numerous specimens obtained in the Shirado-Beds, I became convinced that they are only more fully grown forms of one and the same species. If this has any connection with the sea-temperature, then these two species must be said to thrive best in cooler waters. Anyhow, *Venericardia ferruginea* is now found only in Northern Japan.

Description of the Species

Mollusca

1. *Terebra* sp.

There are several imperfect specimens of a *Terebra* strongly resembling those of *Terebra lischieneana* Dkr. (Yokoyama, Foss. Miura Penin., p. 31, pl. I, fig. 10). The ribs, however, are more prominent with the transverse impressed lines deeper, so that the former appears more or less nodose.

Fossil occurrence.—Kamanomaye (Kadōno)¹⁾.

RCM 22112

2. *Conus* sp.

A mutilated specimen looking much like *Conus sieboldi* Rve (Yokoya-ma, Foss. Miura Penin., p. 34, pl. I, fig. 14).

Fossil occurrence.—North of Toyoma (Toyomamura)²⁾.

RCM 22114

3. *Pleurotoma mediocarinata*, YOKOYAMA.

Pleurotoma mediocarinata, Yokoyama, Foss. Miura Penin., p. 36, pl. I, fig. 18.

Rather common.

Fossil occurrence.—Kagenosaku (Yunami in Sekimoto)³⁾. Lower Musashino.

RCM 22115

4. *Pleurotoma* sp.

Pl. I. Fig. 9.

A specimen with only the last two whorls. It seems to be a large form of *Pleurotoma kamakurana* Pils. (Yokoyama, Foss. Miura Penin., p. 35, pl. I, fig. 17) However, the vertical folds on the body-whorl are

1) 磐城石城郡上遠野釜ノ前 2) 同郡豊間 3) 常陸多賀郡關本村湯網影ノ作

rather weak and indistinct. An accurate determination is at present not possible.

Fossil occurrence.—Izura¹⁾.

5. *Drillia principalis*, PILSBRY.

2CM22116

Drillia principalis. Yokoyama, Foss. Miura Penin, p. 36, pl. I, fig. 20. Foss. Up. Musashino, p. 14, pl. I, fig. 18.

A single example of a young individual, with ribs more prominent than in the adult ones.

Fossil occurrence.—Izura. Lower and Upper Musashino.

Living.—Northern, Central and Western Japan.

6. *Bela rugulata*, Troschel, var. *schniederi*, HARMER.

2CM22117

Bela rugulata var. *schniederi*. Yokoyama, Foss. Up. Musash., p. 44, pl. I, fig. 37.

A single imperfect specimen which, however, undoubtedly belongs to the species above named.

Fossil occurrence.—Enamura Quarry²⁾. Upper Musashino. English Crag.

7. *Olivella fortunei*, (A. ADAMS).

2CM22118

Olivella fortunei. Yokoyama, Foss. Up. Musash., p. 47, pl. II, fig. 3.

2CM22118

A single determinable specimen. Numerous casts from Sukegawa may possibly belong to the same species.

Fossil occurrence.—Futatsujima; Sukegawa³⁾ (Tsurushihama)? Upper Musashino.

17

Living.—Central Japan.

CA122119

8. *Voluta megaspira*, SOWERBY.

CM22120

Voluta megaspira. Yokoyama, Foss. Miura Penin., p. 46, pl. II, fig. 18. Foss. Up. Musashino, p. 50.

(2) CM2212089

Quite frequent. Some specimens are tolerably large.

Fossil occurrence.—North of Toyoma (Toyomamura);⁴⁾ Izura; Sukegawa (Tsurushihama and Hatsuzaki).⁵⁾ Lower and Upper Musashino.

17

Living.—Northern, Central and Western Japan.

1) 五浦 (常陸多賀郡) 2) 江名村 (磐城石城郡) 3) 助川都留志濱 (常陸多賀郡) 4) 豊間村 (磐城石城郡) 5) 初崎

*CM22121-1-**CM22122**CM22123*9. *Chrysodomus phoeniceus*, DALL.

Pl. I. Fig. 1.

Chrysodomus phoeniceus. Foss. Miura Penin., p. 50, pl. I, figs. 8-10.

Several specimens, some of which are quite large.

Fossil occurrence.—Izura; Yunami (Tōzenji)¹⁾; Sukegawa (Tsurushihama). Lower Musashino.

Living.—Japan? British Columbia.

*CM22124-1-2*10. *Chrysodomus pericochlion*, SCHRENCK.

Pl. I. Fig. 2.

*CM22125**CM22126**Chrysodomus pericochlion*. Pilsbry, Cat. Mar. Moll Japan, p. 28.*Tritonium (Buccinum) pericochlion*. Schrenck, Moll. d. nordjap. Meeres. p. 433, pl. xvii, fig. 11, 12.Several specimens, all of which, however, are in the form of casts. In comparing with *Chrysodomus schrencki* Yok. (Foss. Miura Penin., p. 51, pl. III, fig. 1) which may possibly be only a varietal form of Schrenck's species, the whorls are flatter.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki). Lower Musashino.

Living.—Northern Japan.

*CM22127**CM22128**CM22129**CM22130**CM22131**CM22132**CM22133**CM22134*11. *Buccinum leucostoma*, LISCHKE.*Buccinum leucostoma*. Yokoyama, Foss. Up. Musashino, p. 55, pl. II, fig. 11.

Many large examples. Nearly all as casts.

Fossil occurrence.—Oyazaku¹⁾; Sukegawa (Tsurushihama and Hatsuzaki). Upper Musashino.

Living.—Central Japan.

12. *Volutaripa perryi*, (JAY).*Volutaripa perryi*. Yokoyama, Foss. Miura Penin., p. 55, pl. IV, fig. 11. Foss. Up. Musashino, p. 57, pl. II, fig. 19.

This is a very variable species, the young individuals being much more slender than the adult ones which are rather globular. The fossil found in Hitachi, though quite common, belongs to the former.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki). Lower and Upper Musashino.

Living.—Northern and Central Japan.

1) 關本村湯網東禪寺 (多賀郡) 2) 小谷作 (好間村)

Molluscan Remains from the Uppermost Part of the Jō-Ban Coal-Field.

11

CM 22140

CM 22141

13. *Priene oregonensis*, (REDFIELD).

Priene oregonensis. Yokoyama, Foss. Miura Penin., p. 64, pl. III, fig. 10, 12.
Foss. Up. Musashino, p. 68.

Numerous, though mostly found as casts.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki). Lower and Upper Musashino.

Living.—Northern Japan. Alaska, Oregon.

R CM 22142-1-10

CM 22143

14. *Galeodea (Sconsia) japonica*, YOKOYAMA.

Pl. I. Fig. 10.

Galeodea (Sconsia) japonica. Yokoyama, Foss. Moll. Neog. Izumo, p. 3, pl. I,
fig. 4.

A few specimens not well preserved.

Fossil occurrence.—Izura; Yunami (Tōzenji). Pliocene of Izumo.

CM 22144-1-5

CM 22145

Pl. I. Fig. 5.

Cassis japonica. Reeve, Conch. Icon., Cassis, no. 23, pl. IX, fig. 23. Mart.
Chemn. Syst. Conch. Cab., III, part 1, Cassis p. 24, pl. 44, fig. 5. Lischke, Jap.
Meeresconch., I, p. 65. Dunker, Index Moll. Mar. Jap., p. 64.

This shell which is characterized by its swollen oval shape with a short conical spire and convex whorls ornamented with numerous spiral grooves is frequently found as casts.

Fossil occurrence.—Sukegawa (Tsurushihama).

Living.—Central and Western Japan.

R CM 22146

16. *Dolium luteostomum*, KÜSTER.

Dolium luteostomum. Yokoyama, Foss. Miura Penin., p. 66, pl. IV, fig. 2.
Foss. Up. Musashino. p. 69, pl. III, fig. 10.

A small young individual.

Fossil occurrence.—Sukegawa (Tsurushihama). Lower and Upper Musashino.

Living.—Northern, Central and Western Japan. Indian Ocean.

RCM22147-2-6

17. *Cerithium baculum*, YOKOYAMA.

RCM22148

Pl. II. Figs. 6.

Cerithium baculum. Yokoyama, Tert. Foss. Kii, p. 52, pl. VI, fig. 12.

Some time ago, I described from Kii a species of *Cerithium* under the above name. It was founded on a single imperfect specimen. Since then better ones have been found in Iwaki. From these, in spite of their being outer impressions, by taking their casts, we can get a tolerably good idea of their external characters.

The specimen figured is a nearly perfect one, though not the largest. The shell is high-turreted, acute at apex and is composed of about seventeen whorls which are nearly flat, except the last one which is rounded towards the periphery. The ornamentation consists of a subsutural row of spine-like tubercles, some seventeen or eighteen in number on the body-whorl and several (usually six) unequal spiral cords of which the lowest or suprasutural one is the largest. Below the rounded periphery there are about ten spiral cords more or less equal in size. Some of the cords on the whorls show a granular or bead-like aspect. It is about 49 millim. in height and 15 millim. in diameter.

Fossil occurrence.—Yamoto in Kadōno¹⁾; boundary between Kadōno and Iwasaki²⁾. Pliocene of Kii.

RCM22149-2-7

18. *Cerithium* sp.

Pl. II. Fig. 7.

Several impressions of a large *Cerithium*, much larger than the preceding, but having a smaller apieal angle and fewer tubercles, thus approaching a form known as *Vicarya callosa* Jenkins from the Miocene of Java and Philippines and also reported from some parts of Japan.

Fossil occurrence.—Close to Kamanomaye³⁾ and Yamoto, both in Kadōno.

CM22150

19. *Vermetus* sp.

Casts of *Vermetus* resembling *Vermetus tricarinatus* Yokoyama (Moll. Coral-Bed. Awa, p. 24, pl. I, figs. 17–19) in the mode of convolution.

Fossil occurrence.—Sukegawa (Tsurushihama).

1) 矢本（磐城石城郡上遠野） 2) 釜ノ前 3) 鶴崎

Molluscan Remains from the Uppermost Part of the Jō-Ban Coal-Field.

13

CM 22151-2-8

CM 22152

CM 22153

CM 22154

CM 22155

20. *Turritella nipponica*, YOKOYAMA.

Pl. II. Fig. 8.

Turritella nipponica. Yokoyama, Foss. Miura Penin., p. 71, pl. IV, figs. 16-19.

Not rare. Of the four prominent spiral ribs found on each whorl, the three lower ones may be nearly equal in size, leaving the uppermost as the smallest.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki); Yunami (Tozenji and Kagenosaku)¹⁾; Izura; Enamura Quarry; Harakida²⁾. Lower Musashino.

21. *Solarium* sp.

CM 22157

A cast which seems to be that of *Solarium perspectivum* Linné (Tryon, Man. Conch., IX, pl. II, fig. 18-21).

Fossil occurrence.—Sukegawa (Tsurushihama).

CM 22158

CM 22159

CM 22160

22. *Calyptarea mammilaris*, (BRODERIP).

Calyptarea mammilaris. Yokoyama, Foss. Miura Penin., p. 75, pl. IV, fig. 5. Foss. Up. Musashino, p. 82. Moll. Rem. Lowest Part Jō-Ban Coal-field, p. 11, pl. I, fig. 17.

Several large specimens, one of which is about 26 millim. in diameter and 13 millim. in height. In most cases they are deprived of their shell.

Fossil occurrence.—Yunami (Tozenji); Futatsujima³⁾; Tempizan. Lower and Upper Musashino. Asagai-Beds.

Living.—West coast of America from Puget Sound down to the Strait of Magellan.

CM 22161

23. *Crepidula* sp.

Several casts of a large *Crepidula* up to 40 millim. in the greatest diameter. They suggest some of the grown forms of *Crepidula aculeata* (Gm.) (Tryon, Man. Conch., VIII, pl. 39, figs. 61-65) or possibly also *Crepidula grandis* Mid. (Dunker, Index Moll., pl. VI, figs. 1, 2).

CM 22162

CM 22163

CM 22164

CM 22165

CM 22166

Fossil occurrence.—Tempizan.

24. *Natica janthostoma*, DESHAYES.

Natica janthostoma. Yokoyama, Foss. Miura Penin., p. 77, pl. V, figs. 3, 4. Foss. Up. Musashino, p. 83. Foss. Moll. Neog. Izumo, p. 4. Tert. Moll. Dainichi, p. 12. Tert. Foss. Kii, p. 53. Moll. Remains Lowest Part Jō-Ban Coal-field, p. 12, pl. I, fig. 20.

1) 影ノ作 (湯瀬) 2) 原木田 (磐城石城郡玉川村) 3) 二ツ鳴 (磯原)

Frequent.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki); Tempizan¹⁾; Yunami (Tozenji); Izura. Lower and Upper Musashino. Pliocene of Izumo, Dainichi and Kii. Asagai-Beds.

Living.—Northern and Central Japan. Kamchatka.

CM 22167

CM 22168

25. *Polinices (Neverita) ampla*, (PHILIPPI).

Polinices (Neverita) ampla. Yokoyama, Foss. Miura Penin., p. 78, pl. V, figs. 5, 6. Foss. Up. Musash., p. 84. Tert. Moll. Dainichi, p. 12. Tert. Foss. Kii, p. 53.

Frequent.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki). North of Toyoma? Lower and Upper Musashino. Pliocene of Kii and Dainichi.

Living.—Northern to Western Japan, Philippines, etc.

CM 22169

26. *Sigaretus (Eunaticina) papilla*, (GMELIN).

Sigaretus (Eunaticina) papilla. Yokoyama, Foss. Up. Musash., p. 84, pl. V, fig. 8, Tert. Moll. Dainichi, p. 12.

A single example.

Fossil occurrence.—Izura. Upper Musashino.

Living—Central and Western Japan. Philippines and Moluccas.

CM 22170

27. *Turbo (Batillus) cornutus*, GMELIN.

Turbo (Batillus) cornutus. Yokoyama, Moll. Coral-Bed Awa, p. 31, pl. I, fig. 22.

Several specimens, some of which are very large, but all in the form of casts.

Fossil occurrence.—Sukegawa (Tsurushihama). Pleistocene of Awa.

Living.—Northern to Southern Japan. China Sea.

CM 22171-1-3

28. *Bathybembix argenteo-nitens*, LISCHKE.

Pl. I. Fig. 3.

Bathybembix argenteo-nitens. Pilsbry, Catal., p. 97.

Trochus argenteo-nitens. Lischke, Jap. Meeresconch., III, p. 16, pl. IV, fig. 1.

A beautiful shell with the whorls furnished with a single angle, except the last which shows two such angles, the lower one being formed by the periphery. Some specimens are up to 40 millim. in diameter.

Very frequent.

Fossil occurrence.—Sukegawa (Tsurushihama).

Living.—Central and Western Japan.

1) 天妃山 (常陸多賀郡磯原)

29. *Stomatella japonica*, A. ADAMS.

CM 22173 - 2-3

Pl. II. Fig. 3.

Stomatella japonica. A. Adams, Proc. Zool. Soc. London, Feb. 1850, p. 31. Pilsbry, Catal., p. 104; Dunker Moll. Jap., p. 23. Index Moll., p. 145. Sowerby, Thes. Conch., II, p. 838, pl. 174, fig. 14.

This species is allied to *Stomatella lyrata* Pils. (Yokoyama, Foss. Up. Musashino, p. 115, pl. VI, fig. 2) from which it is distinguished by its whorls being more closely ribbed and nodulous.

A single imperfect specimen about 30 millim. in diameter.

Fossil occurrence.—Sukegawa (Tsurushihama).

Living.—Western Japan.

CM 22174

CM 22175

30. *Haliotis* sp.

Very frequent as casts which in all probability belong to *Haliotis diversicolor* Rve. (Pilsbry, Man. Conch., XII, pl. 104, fig. 105) living in Central and Western Japan, and also in China, Australia, etc.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki).

CM 22176

31. *Acmaea* sp.

A single mutilated cast, oval in shape 35 millim. long, 28 millim. broad and about 10 millim. high. It seems to be that of *Acmaea nigro-lineata* Rve. (Lischke, Jap. Meeres conch., vol. I, pl. VIII, figs. 5-11), living in our seas.

Fossil occurrence.—Tempizan.

(R) CM 22177 - 1-4

(3) CM 22178

CM 22179

X CM 22180

32. *Helcioniscus pallidus*, (GOULD).

Pl. I. Fig. 4.

Helcioniscus pallidus. Yokoyama, Foss. Miura Penin., p. 101, pl. VI, figs. 16, 17. Foss. Up. Musash., p. 117.

This is a very frequent shell, though mostly as casts.

Fossil occurrence.—Yunami (Tozenji); Sukegawa (Tsurushihama and Hatsuzaki). Lower and Upper Musashino.

Living.—Northern and Central Japan.

R CM 22181

33. *Dentalium weinkauffi*, DANKER.

Dentalium weinkauffi. Yokoyama, Foss. Miura Penin., p. 102, pl. VI, figs. 19-21.
Foss. up. Musashi., p. 118, pl. VI, fig. 6.

Mostly present as casts.

Fossil occurrence.—Sukegawa (Tsurushihama). Lower and Upper Musashino.

Living.—Central Japan.

R CM 22182

R CM 22183

R CM 22184

R CM 22185

34. *Dentalium complexum*, DALL.

Pl. II. Fig. 9.

Dentalium complexum. Yokoyama, Foss. Miura Penin., p. 101, pl. VI, fig. 27.
Foss. Moll. Neog. Izumo, p. 4.

Several specimens, but differing from the preceding by its more rapidly tapering shape toward the apical end.

Fossil occurrence.—Tempizan; between Nijida and Shimoyamada¹⁾; Nonohide²⁾ in Kadono; Toyoma. Lower and Upper Musashino. Pliocene of Izumo.

Living.—Central Japan, Sandwich Islands.

35. *Panope generosa*, (GOULD).

Pl. VI. Fig. 6.

Panope generosa. Yokoyama, Foss. Up. Musash., p. 121. pl. V, figs. 14, 15.
Foss. Moll. Neog. Izumo, p. 4. Tert. Moll. Dainichi, p. 14.

Fossil occurrence.—Yunami (Tōzenji); Harakida; between Nijida and Shimoyamada; north of a triangulation-point in Shimoyamada; Iida³⁾ and Misawa (Kashima); Kamanomaye in Kadono; railway-cutting, Nakoso⁴⁾. Pliocene of Izumo and Dainichi.

Living.—Northern Japan. West coast of North America.

R CM 22194

36. *Mya arenaria* Linne, var. *japonica*, JAY.

Pl. VI. Fig. 4.

Mya arenaria var. *japonica*. Pilisby, Catal., p. 118.

Mya arenaria. Lischke, Jap. Meeresconch., I, p. 138.

Mya japonica. Jay, Rep. Shells Jap. Explor. Exped. Perry, p. 292, pl. I, figs. 7,

10.

Two specimens somewhat imperfect. The Japanese form is generally considered as a variety of the typical one, on account of the shell being more pointed behind. Our fossil unfortunately lacks the posterior end.

1) 仁井田・下山田間 2) 野々日堂 3) 廣島村飯田・三澤 4) 勿來

Fossil occurrence.—Yunami (Tōzenji).

Living.—Northern, Central and Western Japan.

37. *Mactra sulcataria*, DESHAYES.

CM 22195

CM 22196

CM 22197

Mactra sulcataria. Yokoyama, Foss. Up. Musash., p. 126, pl. VII, fig. 6.

Two casts, one of which seems somewhat longer than usual.

Fossil occurrence.—Sukegawa (Tsurushihama); Tempizan. Upper Musashino.

Living.—Northern, Central and Western Japan. North China. Possiet Bay.

CM 22198

38. *Mactra veneriformis*, DESHAYES.

Mactra veneriformis. Yokoyama, Foss. Miura Penin., p. 109, pl. VIII, fig. 10. Foss. Up. Musash., p. 127.

A few isolated valves.

Fossil occurrence.—Yunami (Tōzenji). Lower and Upper Musashino.

Living.—Northern, Central and Western Japan.

CM 22198-1-7

CM 22199

CM 22200

CM 22201-1-8

39. *Mactra spectabilis*, LISCHKE.

Pl. I. Figs. 7, 8.

Mactra spectabilis. Lischke, Jap. Meeresconch., II, p. 120, pl. XI. figs. 1, 2. CM 22202 ②
Pilsbry, Cat., p. 119.

This is a large and beautiful species which is distinguished from *Mactra sulcataria* by its more inequilateral and triangular shape as well as by its larger growth. The beaks are also more pointed and bent. Mostly present as casts.

Fossil occurrence.—Izura; Nakayama¹⁾; Nakanosaku²⁾; Tsuchibashi in Kadōno; Aoya in Watanabe³⁾.

Living.—Central and Western Japan.

CM 22203-2-11

CM 22204

CM 22205

40. *Spisula grayana*, (SCHRENCK).

Pl. II. Fig. 11.

Spisula grayana. Yokoyama, Foss. Up. Musash., p. 130, pl. VIII, figs. 1, 2.

Several casts.

Fossil occurrence.—Sukegawa (Tsurushihama); Tempizan; Fujiwara (opposite the river).

Living.—Northern Japan. Okhotsk Sea. Bering Sea.

1) 石城郡飯野村中山 2) 同江名村中ノ作 3) 同渡邊村青谷

CM 22206

18

Art. 5.—M. Yokoyama:

41. *Solen grandis*, DUNKER.

Solen grandis. Yokoyama, Foss. Up. Musash., p. 134, pl. IX, fig. 4.

Only fragments which are, however, readily recognized by the large size of the shell.

Fossil occurrence.—Tempizan. Upper Musashino.

Living.—Western Japan. Philippines.

CM 22207

42. *Solen gouldii*, CONRAD.

Pl. II. Fig. 5.

Solen gouldii. Conrad, Amer. Jour. Conch., III, Appendix, p. 28. Dunker, Ind. Moll., p. 178, pl. XVI, fig. 11.

Solen gracilis. Gould (non Phil.), Proc. Bost. Soc. Nat. Hist., VIII, p. 26. Otia Conch., p. 165. Lischke, Jap. Meeressconch., II, p. 123.

Two isolated valves. More slender than *Solen krusensterni* Schr. (Foss. Up. Musash., pl. IX, fig. 5) so frequent in the Upper Musashino in the neighbourhood of Tokyo.

Fossil occurrence.—Yunami (Tozenji).

Living.—Northern to Central Japan.

CM 22208

CM 22209

CM 22210-5-3

CM 22211

CM 22212

CM 22213-5-2

CM 22214

CM 22215

CM 22216

43. *Cultellus izumoensis*, YOKOYAMA.

Pl. V. Figs. 2, 3.

Cultellus izumoensis. Yokoyama, Foss. Moll. Neog. Izumo, p. 5, Pl. II, fig. 1.

When I first described this species, the only specimen at hand was broken in front. Now we have several and much larger ones which have the anterior end perfect. It is rounded as was previously presumed, but much more sharply than the posterior end.

Quite frequent.

Fossil occurrence.—North of a triangulation-point, Shimoyamada; between Niida and Shimoyamada¹⁾; Iida in Kashima²⁾; Enamura Quarry³⁾; Ushiroda in Uyeda⁴⁾; Coast of Nakoso. Pliocene of Izumo.

44. *Siliqua pulchella*, (DUNKER).

Siliqua pulchella. Yokoyama, Foss. Up. Musashino, p. 135, pl. IX, fig. 7.

Almost always as casts.

Fossil occurrence.—North of Kamikamado and also Aoya, Watanabe; east of Kamitakaku in Iinomura⁵⁾; near the railway station of Taira. Upper Musashino.

Living.—Central and Western Japan.

1) 仁井田下山田間 2) 鹿鳴村飯田 3) 江名村石切場 4) 植田後田 5) 飯野村上高久

Molluscan Remains from the Uppermost Part of the Jō-Ban Coal-Field.

19

45. *Soletellina violacea*, LAMARCK.

2 CM 22217
2 CM 22218
2 CM 22219

Soletellina violacea. Yokoyama, Foss. Up. Musashino, p. 187, pl. IX, figs. 13, 14.

Rather frequent at some places.

Fossil occurrence.—Kamanomaye and Yamoto in Kadōnō; Tabe in Izumi¹⁾; Kawakamimaye in Iwasaki.²⁾

Living.—Northern to Western Japan. Philippines. Moluccas.

46. *Tellina besshoensis*, YOKOYAMA.

2 CM 22220

Tellina besshoensis. Yokoyama, Moll. Rem. Lowest Part. Jō-Ban Coal-Field, p. 14, pl. II, figs. 1-5.

Several specimens.

Fossil occurrence.—Kamanomaye in Kadōnō. Asagai-Beds (Miocene).

2 CM 22221
2 CM 22222
2 CM 22223

47. *Tellina optiva*, YOKOYAMA.

Tellina optiva. Yokoyama, Foss. Moll. Neog. Izumo, p. 6, pl. II, figs. 3, 4, 2 CM 22224

Frequent.

Fossil occurrence.—Near Takakuraji³⁾; north of a triangulation-point, Shimoyamada; Nakoso-coast. Ushiroda (Uyeda); Ōishi (Fujiwara, Yumoto).⁴⁾ Pliocene of Izumo.

2 CM 27090
2 CM 27091

2 CM 22224-2-12

48. *Tellina izurensis*, YOKOYAMA.

Pl. II. Fig. 12.

There is only a left valve which is to a greater part deprived of its shell, but so characteristic in outline that there is none which can be confounded with it, either among the fossil or living forms hitherto found in Japan.

It is about 52 millim. long and 37 millim. high, roughly triangular in outline, compressed, almost equilateral, the posterior half being only a little longer than the anterior. The anterior border together with the antero-dorsal forms one broadly rounded arch. The posterior border is also rounded, but much more sharply, the postero-dorsal sloping down at a tolerably high angle and insensibly passing into it. The ventral border is straight except at both ends where it curves to meet the anterior and posterior borders respectively, without, however, making any sharp angle. The surface of the shell seems to have been smooth, as can be judged from a little of it still preserved.

Fossil occurrence—Izura.

1) 石城郡泉村川部 2) 磐崎村川上前 3) 高倉寺 4) 湯本町藤原大石

CM22225
RM CM22226

49. *Macoma praetexta*, (MARTENS).

Macoma praetexta. Yokoyama, Foss. Up. Musashino, p. 142, pl. X, figs. 2, 3.
Moll. Remains Lowest Part Jō-Ban Coal-Field, p. 13.

Rare.

Fossil occurrence.—Sukegawa (Tsurushihama). Kokozura near Hirakata. Upper Musashino. Asagai-Beds (Miocene).
Living.—Central and Western Japan.

RM CM22227
RM CM22228
RM CM22229
RM CM22230
RM CM22231-5-9
RM CM22235

50. *Macoma dissimilis*, (MARTENS).

Pl. V. Fig. 9.

Macoma dissimilis. Yokoyama, Foss. Miura Penin., p. 116, pl. VII, figs. 19, 20.
Foss. Up. Musash., p. 143, pl. X, fig. 4.

A few undoubted examples and several doubtful casts.

Fossil occurrence.—Nakayama; Toyoma; Futatsujima; Sukegawa (Tsurushihama); Izura? Tempizan?

Living.—Central Japan.

RM CM22232
RM CM22233
RM CM22234
RM CM22235

51. *Dosinia troscheli*, LISCHKE.

Dosinia troscheli. Yokoyama, Foss. Miura Penin., p. 119, pl. VIII, figs. 5, 6.
Foss. Up. Musashino, p. 144. Tert. Moll. Dainichi, p. 15.

Rare.

Fossil occurrence.—Futatsujima; Misawa²⁹; Tempizan; Sukegawa (Hatsuzaki). Lower and Upper Musashino. Pliocene of Dainichi.

Living.—Central and Western Japan.

RM CM22236

52. *Cyclina chinensis*, (CHEMNITZ).

Cyclina chinensis. Yokoyama, Foss. Miura Penin., p. 119, pl. XI, figs. 7, 8.
Foss. Up. Musashino, p. 145.

Fossil occurrence.—Kamanomaye in Kadōno. Lower and Upper Musashino.

Living.—Northern, Central and Western Japan. China Sea. Cochin China.

RM CM22237
RM CM22238
RM CM22239
RM CM22240
RM CM22241-3-2
RM CM22242-3

53. *Meretrix iizukai*, YOKOYAMA.

Pl. III. Figs. 2, 3.

Shell moderate in size and thickness, convex, obliquely oval, very inequilateral, anterior side very short, rounded in front and behind,

RM CM22243
CM22244

1) 石城郡庵田村九面(平潟港附近) 2) 三澤

though more sharply in the former, ventral border broadly arched. Surface only with rude lines of growth. Beaks nearly touching, swollen, incurved, pointed. A cordate lunula is present usually bounded by a shallow line on both sides. Sinus shallow, triangular, usually blunt at end and somewhat directed upward. The length being taken as 10, the height is about 8 and the thickness about 6 on an average. The dentition is at present unknown. The largest example is 73 millim. long, 56 millim. high and 46 millim. thick.

The outward shape is remarkably like that of *Saxidomus purpuratus* Sow. (Yokoyama, Foss. Miura Penin., pl. IX, fig. 8. Foss. Up. Musashino, pl. XII, fig. 9) living and fossil in Japan. But by a careful comparison with the latter, it is somewhat thinner, more inequilateral, with the pallial sinus shorter and more triangular and usually with a lunula more or less bounded by shallow depressions on both sides which is not present in *Saxidomus purpuratus* of about equal size. But it may possibly turn out to be only a variety of the latter which, however, must remain undecided as long as the dentition is not known.

This shell is also not unlike in shape *Meretrix lamarcki* Agassiz of the Miocene of Europe (Cossman et Perot, Conch. Neog. Aquit., pl. 15, figs. 11-15. Höernes, Foss. Moll. Wiener Becken, pl. 18, fig. 5).

The shell seems to be variable in shape, as some of the example have the beaks more swollen than in the others. Rather frequent.

Fossil occurrence.—Between Niida and Shimoyamada; Takakuraji; Harakida; Kekozura near Hirakata; Izura; Yunamé (Tōzenji); Sukegawa (Tsurushihama and Hatsuzaki).

54. *Clementia speciosa*, YOKOYAMA.

CM 22245
CM 22246-1-6

Pl. I. Fig. 6.

Clementia speciosa. Yokoyama, Tert. Moll. Dainichi, p. 15, pl. II, figs. 14, 15.

A few young specimens.

Fossil occurrence.—Between Niida and Shimoyamada; (Railway cutting, Nakoso. Pliocene of Dainichi.

CM 22247

55. *Venus (Mercenaria) stimpsoni*, GOULD.

Venus (Mercenaria) stimpsoni. Yokoyama, Foss. Up. Musashino, p. 148, pl. XI, figs. 11, 12. Foss. Moll. Izumo, p. 6, pl. I, fig. 5.

A very young individual.

Fossil occurrence.—Izura. Upper Musashino. Pliocene of Izumo.

Living.—Northern to Western Japan.

CM 22248

56. *Venus jedoensis*, LISCHKE.

Venus jedoensis. Yokoyama, Foss. Miura Penin., p. 120, pl. VIII, figs. 9, 10. Foss. Up. Musash., p. 148. Moll. Coral-bed Awa, p. 43, pl. II, fig. 20.

A few casts.

Fossil occurrence.—Sukegawa (Hatsuzaki). Lower and Upper Musashino. Pleistocene of Awa.

Living.—Central and Western Japan.

CM 22249

CM 22250

57. *Venus toreuma*, GOULD.

Venus toreuma. Yokoyama, Moll. Coral-Bed Awa, p. 44. pl. II, fig. 22.

Venus crebrisulea. Sowerby, Thes. Conch., II, p. 728, pl. 161, figs. 187-189.

Venus jukesii. Syst. Conch., Cab. Mart. Chem., Venus, p. 217, pl. 35, figs. 7-9.

This is a thick-shelled, swollen, almost globular shell with raised, lamellar, concentric ribs. Rare.

Fossil occurrence.—Tempizan; Sukegawa (Hatsuzaki). Pleistocene of Awa.

Living.—Central and Western Japan. Philippines. Australia.

CM 22251-57

CM 22252

58. *Tapes undulatus*, (BORN).

Pl. V. Fig. 1.

Tapes undulatus. Dunker, Index Moll., p. 206. Pilsbry, Catal., p. 129.

Venus undulata. Born, Mus. Caes. Vind., p. 67.

Venus rimosa. Philippi, Abbild., III, pl. VII, fig. 7.

A transversely much elongated shell and rather compressed, with both ends rounded. The proportion of length to height is about 10 to 6. A few casts.

Fossil occurrence.—Sukegawa (Hatsuzaki).

Living.—Central and Western Japan.

CM 22253

59. *Cardium californiense*, DESHAYES.

Cardium californiense. Yokoyama, Foss. Miura Penin., p. 127, pl. IX, fig. 10. Foss. Up. Musashino, p. 154.

A single cast.

Fossil occurrence.—Tempizan. Lower and Upper Musashino.

Living.—Northern to Western Japan. Behring Sea to California.

CM 22254

60. *Cardium muticum*, REEVE.

Cardium muticum. Yokoyama, Foss. Miura Penin., p. 128, pl. IX, fig. 11.
Foss. Up. Musash., p. 155.

A large and good example.

Fossil occurrence.—Komeda in Kashima¹. Lower and Upper Musashino.

Living.—Northern, Central and Western Japan. Philippines. East Indies.

CM 22255

61. *Cardium modestum*, ADAMS ET REEVE.

Cardium modestum. Yokoyama, Foss. Miura Penin., p. 128, pl. IX, figs. 12, 13.
Foss. Up. Musash., p. 55.

Numerous casts which somewhat look like those of the next species, but readily recognized by the impression of much finer ribs at the ventral border.

Fossil occurrence.—Sukegawa (Hatsuzaki).

Living.—Central Japan.

CM 22256
CM 22257
CM 22258
CM 22259
CM 22260
CM 22261

62. *Cardium shinjiense*, YOKOYAMA.

Cardium shinjiense. Yokoyama, Foss. Moll. Neog. Izumo, p. 7, pl. II, fig. 6.
Moll. Rem. Lowest Part. Jō-Ban Coal-Field, p. 16, pl. III, figs. 13-15.

Very frequent, but mostly as casts and more or less distorted.

Fossil occurrence.—Sukegawa (Tsurushihama); Tempizan; Yunami (Tōzenji and Kagenosaku); Ushiroda, Uyeda; Kamanomaye in Kadono; north of the triangulation-point in Shimoyamada. Pliocene of Izumo. Iwaki- and Asagai-Beds.

CM 22262
CM 22263

63. *Diplodonta semiaspera*, PHILIPPI.

Diplodonta semiaspera. Yokoyama, Foss. Miura Penin., p. 160, pl. XIV, fig. 2.
Foss. Up. Musash., p. 180, pl. X, figs. 2, 3.

Many casts.

Fossil occurrence.—Kamikamado in Watanabe²; Sukegawa (Hatsuzaki). Lower and Upper Musashino.

Living.—Central and Western Japan. West Indies.

1) 鹿嶋村米田 2) 渡邊村上釜戸

CM 22264
 CM 22265
 CM 22266
 CM 22267
 CM 22268
 CM 22269
 CM 22270-5-5
 CM 22271
 CM 22272
 CM 22273
 CM 22274-5-6
 CM 22275-7
 CM 22276-8
 CM 22277
 CM 22278

64. *Lucina (Phacoides) borealis*, (LINNE).
 Pl. V. Figs. 5-8.
Lucina borealis. Yokoyama, Foss. Miura Penin., p. 133, pl. X, fig. 7. Foss.
 Up. Musashino, p. 160, Tert. Foss. Kii, p. 67, pl. VI, fig. 11.
Lucina spectabilis. Yokoyama, Foss. Miura Penin., p. 134, pl. X, figs. 10-12.
 Very frequent, some being very large. *Lucina spectabilis* which I described from the Lower Musashino of Koshiba represents, as I am now convinced, only very large forms of the same species. The shell is variable in shape. The living ones are small.
 Fossil occurrence.—Taira (Park); Komeda in Kashima; between Kinuya and Yakuōji¹⁾; Toyoma; Enamura Quarry; Iwasaki; Sukegawa (Tsurushihama and Hatsuzaki); Tempizan; Yunamii (Tōzenji); Izura, Lower and Upper Musashino. Pliocene of Kii. Miocene, Pliocene and Pleistocene of Europe.

CM 22279
 CM 22280
 CM 22281
 CM 22282-6-5
 CM 22283
 CM 22284

Living.—Central Japan. Atlantic and Mediterranean Sea.

65. *Thyasira bisecta*, (CONRAD).

Pl. VI. Fig. 5.

Thyasira bisecta. Yokoyama, Moll. Rem. Lowest Part Jō-Ban Coal-Field, p. 18,
 pl. III, fig. 2.
 Several examples better preserved than that figured in the above work.
 Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki); Ishiwaki in Kabeya²⁾; Taira (Park); Harakida; Iwasaki (Kawakamimaye). Asagai-Beds. Miocene and Pliocene of the Pacific coast of North America.

CM 22285
 CM 22286

Living.—Coast of Alaska. Puget Sound.

66. *Cardita cumingiana*, DUNKER.

Cardita cumingiana. Yokoyama, Foss. Miura Penin., p. 137, pl. X, fig. 16,
 pl. XI, fig. 1.

Rare.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki). Lower Musashino.

Living.—Central and Western Japan.

67. *Venericardia ferruginea*, (ADAMS).

Pl. V. Fig. 4.

Venericardia ferruginea. Yokoyama, Foss. Miura Penin., p. 139, pl. XI, fig. 3, 4.
 Foss. Up. Musashino, p. 162.

Venericardia sp. Yokoyama, Moll. Rem. Lowest Part Jō-Ban Coalf., p. 19,
 pl. III, figs. 8, 9.

1) 草野村綱谷、大野村薬王子ノ間 2) 神谷村石脇

The specimens attain a much larger size than those hitherto found fossil or living. The length attained is 30 millim., that is to say, it is one and a half times larger.

A shell which I described from the Iwaki-Beds under the name of *Venericardia* sp. belongs, as I am now convinced, to this same species.

Fossil occurrence.—Harakida; Komeda in Kashima; Toyoma; Enamura Quarry; Kokozura near Hirakata; Tsuchibashi and Nonohido in Kadono¹⁾; Yūnami (Tōzenji). Lower and Upper Musashino. Iwaki-Beds.

Living.—Northern Japan.

R CM 22295-111

R CM 22296

68. *Thracia pubescens*, PULTENEY.

Pl. I. Fig. 11.

Thracia pubescens. Yokoyama, Foss. Shells Saishu, p. 6, pl. I, fig. 1.

A few, though undoubted specimens.

Fossil occurrence.—Gomasawa in Taira.²⁾ Upper Musashino of Saishu. Crag of England.

Living.—Atlantic.

R CM 22297-2-10

R CM 22298

69. *Mytilus grayanus*, DUNKER.

Pl. II. Fig. 10.

Mytilus grayanus. Dunker, Zeitsch. f. Malac., 1853, p. 84. Index Moll., p. 221. Lischke, Jap. Meeressch., I, p. 153, pl. X, figs. 7, 8.

Mytilus dunkeri. Reeve, Conch. Icon., Mytilus, sp. 17.

Several casts of this beautiful triangular compressed shell.

Fossil occurrence.—Tempizan.

Living.—Northern Japan. Sakhalin, Philippines, Tasmania, etc.

R CM 22299

70. *Mytilus* sp.

A single cast, smaller, more elongated and inflated, looking that of *Mytilus hirsutus* Lam. fossil and living in Japan.

Fossil occurrence.—Tempizan.

X CM 22300

71. *Modiola* sp.

A cast very much like that of *Modiola barbata* L., fossil and living in Japan.

Fossil occurrence.—Sukegawa (Tsurushihama).

1) 上野野橋及七野々日堂 2) 平町ガマ澤

CM 22307

Art. 5.—M. Yokoyama:

72. *Lima lima*, (LINNÉ).*Lima lima*. Yokoyama, Moll. Coral-Bed Awa, p. 54, pl. III, fig. 13.

A single cast.

Fossil occurrence.—Sukegawa (Tsurushihama). Pleistocene of Awa. Pliocene of Italy. Miocene of Vienna.

Living.—Central to Southern Japan. Red Sea. Mediterranean Sea.

CM 22302

73. *Lima angulata*, SOWERBY.*Lima angulata*. Yokoyama, Foss. Miura Penin., p. 148, pl. XII, fig. 12. Foss. Up. Musash., p. 177.

Two small isolated valves.

Fossil occurrence.—Sukegawa (Tsurushihama). Lower and Upper Musashino. Neogene of New Zealand.

Living.—Northern and Central Japan. Philippines. Australia.

CM 22303

74. *Lima vulgatula*, YOKOYAMA.*Lima vulgatula*. Yokoyama, Foss. Up. Musashino, p. 179, pl. XVII, figs. 18, 19.

A single valve, somewhat larger than those of the Upper Musashino and also with the ribs a little more numerous.

Fossil occurrence.—Sukegawa (Tsurushihama). Upper Musashino.

R CM 22304-3/

R CM 22305

R CM 22306-2/

CM 22307

75. *Lima goliath*, SOWERBY.

Pl. III.—figs. 1, 4.

Lima goliath. Yokoyama, Foss. Miura Penin., p. 147, pl. XVI, figs. 7, 8.

Two nearly perfect valves with several fragments, one of which is very large.

Fossil occurrence.—Kamanomaye in Kadōno; Tempizan; Sukegawa (Tsurushihama and Hatsuzaki). Lower Musashino.

Living.—Central Japan (Sagami Bay). Southeastern Japan at 750 fathoms (Challenger). Patagonia.

CM 22308

X CM 22309

76. *Pecten laetus*, GOULD.*Pecten laetus*. Yokoyama, Foss. Miura Penin., p. 152, pl. XIV. figs. 1, 2. Foss. Up. Musashino, p. 180, pl. XIV. fig. 20. Foss. Shells Saishu, p. 7. Moll. Coral-Bed Awa, p. 56.

Though not frequent, some specimens are very large.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki); Tempizan. Lower and Upper Musashino. Pleistocene of Awa.

Living.—Northern to Western Japan.

CM22310

77. *Pecten vesiculosus*, DUNKER.

Pecten vesiculosus. Yokoyama, Foss. Miura Penin., p. 154, pl. XIII, figs. 11-13.
Foss. Up. Musashino, p. 181.

A right valve about 40 millim. in length and height.

Fossil occurrence.—Sukegawa (Tsurushihama). Lower and Upper Musashino.

Living.—Central Japan.

78. *Pecten swiftii*, BERNARDI.

CM22311-2-1

Pl. II. Fig. 1.

Pecten swiftii. Yokoyama, Foss. Miura Penin., p. 154, pl. XIV, fig. 11.

A large right valve.

Fossil occurrence.—Sukegawa (Tsurushihama). Lower Musashino.

Living.—Northern Japan. Sea of Okhotsk. Alaska.

12 CM22312
2 CM22313
2 CM22314

79. *Pecten yessoensis*, JAY.

Pecten yessoensis. Yokoyama, Foss. Miura Penin., p. 159, pl. XIII, figs. 14, 15.

Pretty frequent.

Fossil occurrence.—Izura; Tempizan; Isohara. Lower Musashino.

Living.—Northern Japan. Sea of Okhotsk.

2 CM22315-4-4
2 CM22316-4-1
2 CM22317-4-2
2 CM22318-4-3
2 CM22319-4-5

80. *Pecten kimurai*, YOKOYAMA.

Pl. IV. Pl. II. Fig. 4.

Shell large, moderately thick, orbicular, compressed, inequivalve, ~~CM22320-4-6~~ nearly equilateral. Surface with radiating ribs; ribs on the right valve ~~CM22321-4-4~~ usually nine in number, broad, rounded, with a few longitudinal riblets ~~CM22322~~ striae on the back, separated by shallow valleys of about equal breadth ~~CM22323~~ which are smooth at bottom and show no sharp demarcation against the ~~CM22324~~ ribs; ribs on the left valve also usually nine in number, roof-like, sharp ~~CM22325~~ with valleys equally formed, that is to say broadly V-shaped. Ears unequal, the anterior one coarsely and radiately corded, with a shallow rounded ~~CM22326~~ byssal notch below, while the posterior one is equally long, but usually ~~CM22327~~ smooth. Height slightly greater than length; thickness about one-fourth ~~CM22328~~ the height.

The largest specimen measures about 100 millim. in height.

This shell is much like *Pecten tokyoensis* Tok. (Yokoyama, Miura Penin., pl. XIV, fig. 7, 8), but the ribs are more prominent and elevated.

Fossil occurrence.—Izura; Yunami (Tōzenji); Sukegawa (Hatsuzaki); Kashima (Iida and Komeda); Taira (Gomasawa and also near the railway station); Toyoma; Kamitakaku in Iino; Enamura Quarry; Kusano (between Mizushima and Yamazaki)¹⁾; Tamaye, Kobisa,²⁾

R CM 22330

81. *Ostrea gigas*, THUNBERG.

Ostrea gigas. Yokoyama, Foss. Miura Penin., p. 162, pl. XV, figs. 1, 2. Foss. Up. Musashino, p. 184. Moll. Coral-Bed Awa, p. 57.

A single fragment of the beak-portion of a lower valve.

Fossil occurrence.—Tempizan. Lower and Upper Musashino. Pleistocene of Awa.

Living.—Northern, Central and Western Japan. China.

R CM 22331-6-7

82. *Pinna japonica*, HANLEY.

Pl. VI. Fig. 7.

Pinna japonica. Yokoyama, Foss. Up. Musashino, p. 185, pl. XV, fig. 8.

A fragment.

Fossil occurrence.—Iida in Kashima. Upper Musashino.

Living.—Central and Western Japan.

C 1422332

R CM 22333

83. *Arca kobeltiana*, PILSBRY.

Arca kobeltiana. Yokoyama, Foss. Miura Penin., p. 163, pl. XVII, fig. 4. Foss. Up. Musash., p. 185. Moll. Coral-Bed Awa, p. 59.

Quite frequent.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki). Tempizan. Lower and Upper Musashino. Pleistocene of Awa.

Living.—Northern and Central Japan.

R CM 22334

R CM 22335

R CM 22336

R CM 22337

CM 22338

84. *Arca (Scapharca) subcrenata*, LISCHKE.

Arca (Scapharca) subcrenata. Yokoyama, Foss. Up. Musashino, p. 187, pl. XV, fig. 12.

Several imperfect examples.

Fossil occurrence.—Yamamoto (entrance of the coal-mine of Fujiwara); between Niida and Shimo-yamada; Toyoma; Yunami (Tozenji); Sukegawa (Tsurushihama). Upper Musashino.

Living.—Central, Western and Southern Japan.

1) 草野(水晶・山崎ノ間) 2) 小久町田前

85. *Arca setoensis*, YOKOYAMA.

CM 22339
CM 22340
CM 22341

Area setoensis. Yokoyama, Tert. Foss. Kii, p. 58, pl. VII, fig. 13.

A few isolated valves, larger than the specimens described from Kii.

Fossil occurrence.—Yūnami (Tōzenji); between Mizushima and Yamazaki in Kutsanb; Shimanoyamada (north of a triangulation-point). Pliocene of Kii.

86. *Pectunculus vestitus*, DUNKER.

CM 22342

Pectunculus restitus. Yokoyama, Foss. Miura Penin., p. 167, pl. XVII, figs. 10, 11. Foss. Up. Musashino, p. 189, pl. XVI, figs. 1-3. Foss. Shells Saishu, p. 7. Moll. Rem. Lowest Part., p. 21.

Numerous examples.

Fossil occurrence.—Tempizan. Iwaki-Beds. Musashinos.

Living.—Central Japan.

CM 22343
CM 22344

87. *Parallelodon obliquatus*, YOKOYAMA.

Parallelodon obliquatus. Yokoyama, Foss. Miura Penin., p. 170, pl. XVIII, figs. 9-11. Foss. Up. Musash., p. 191. Moll. Coral-Bed Awa, p. 62.

Not rare.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki). Tempizan. Lower and Upper Musashino. Pleistocene of Awa.

Living.—Northern and Central Japan.

X CM 22345

88. *Limopsis azumana*, YOKOYAMA.

Limopsis azumana. Yokoyama, Foss. Miura Penin., p. 174, pl. XVIII, figs. 19-21. Foss. Up. Musashi, p. 193.

Several specimens.

Fossil occurrence.—Sukegawa (Hatsuzaki). Lower and Upper Musashino.

CM 22346
CM 22347
CM 22348
CM 22349

89. *Leda confusa*, HANLEY.

Leda confusa. Yokoyama, Foss. Up. Musash., p. 195. pl. XVII, fig. 4.

Several isolated valves.

Fossil occurrence.—Harakida; Kamitakaku in Iino; Yūnami (Tōzenji); Izura? Upper Musashino.

Living.—Central Japan. China Sea.

CM 22350
R CM 22351

90. *Leda* sp.

Casts of a more elongated form like *Leda ramsayi* Sm. and *Leda gordonis* Yok. All being broken at the posterior end, it is not possible to determine them.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki); Futatsujima.

CM 22352 - 2-2
CM 22353

91. *Yoldia lischkei*, SMITH.

Pl. II. Fig. 2.

Yoldia lischkei. Smith, Challenger Lamellibranchiata, p. 247, pl. XX, fig. 4.

The shell is thin, moderately convex, transversely squarely oblong, somewhat inequilateral, anterior border rounded, posterior subtruncate. Surface smooth. A faint ridge runs from the beak to the postero-ventral corner behind which the surface is a little depressed.

Most of the specimens are found as casts.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki); Yūnami (Tōzenji and Kagenosaku).

Living.—Central Japan (Sea of Sagami at the depth of 345 fathoms).

R CM 22354

92. *Nucula insignis*, GOULD.

Nucula insignis. Yokoyama, Foss. Miura Penin., p. 179, pl. XIX, figs. 7, 8. Foss. Up. Musashi., p. 198. Foss. Shells Saishu, p. 7. Moll. Rem. Lowest Part Jo-Ban Coal-Field, p. 21.

A few examples.

Fossil occurrence.—Sukegawa (Tsurushihama). Lower and Upper Musashino. Asagai-Beds.

Living.—Northern Japan.

CM 22355

93. *Nucula mirabilis*, ADAMS ET REEVE.

Nucula mirabilis. Yokoyama, Foss. Miura Penin., p. 180, pl. XIX, fig. 9. Foss. Shells Neog. Izumo, p. 9.

Several examples.

Fossil occurrence.—Sukegawa (Hatsuzaki). Lower Musashino. Pliocene of Izumo.

Living.—Central and Western Japan.

94. *Solemya tokunagai*, YOKOYAMA.

Pl. VI. Figs. 1, 2, 3.

Several specimens of this interesting shell were collected at three places. Though more or less imperfect, they still give a fairly good idea of its general characters.

The shell is moderate in thickness, compressed, very inequilateral, transversely elongated, more or less rectangular, though broadest in front. The anterior border is truncate, the posterior sharply rounded. The dorsal and ventral borders are almost parallel, the former very slightly arched and the latter somewhat excavated. The antero-ventral corner is rounded. The surface is radiately channelled, with the channels slightly curved in the anterior part, broad, shallow, equal to, or somewhat broader than interspaces, anteriorly distinct, but gradually getting shallower, narrower and indistinct posteriorly, although we can trace them even near the posterior end of the shell. In general the most distinct are the anterior six or seven. The dark periostracum of the channels are elongated into long processes.

The most perfect specimen is that given in fig. 2. It has both valves preserved, only with the postero-ventral corner broken. It is 94 millim. long and 38 millim. high. Another with the processes of the anterior part preserved (fig. 1) is broken at the posterior end. It is somewhat longer than the other, being presumably over 70 millim. long with the height only 22 millim. It may possibly belong to a different species, but the general outline being quite similar, at present I can, at the utmost, only treat it as a variety.

Solemya johnsoni Dall (Moll. and Brach. dredged in Deep Water, Proc. U.S. Nat. Museum, Vol. XVII, pl. XXV, fig. 1) and *Solemya agassizii* Dall (Moll. a. Brach., Rep. Dredg. Oper. W. Coast of America, pl. XVI, fig. 10) of the Western Pacific may be compared to the Japanese fossil, without being perfectly identical.

Fossil occurrence.—Shiogu in Tatsuta; Tsuchibashi in Kadōno; Engamura Quarry.

Brachiopoda.

95. *Terebratulina caput-serpentis*, (LINNÉ).

Terebratulina caput-serpentis. Yokoyama, Foss. Miura Penin., p. 182, pl. XIX, figs. 15-18. Hayasaka, Some Tert. Brach. Japan, p. 7, pl. I, figs. 9-12.

1) 龍田村鹽具

Only as casts.

Fossil occurrence.—Sukegawa (Tsurushihama and Hatsuzaki). Lower Musashino. Miyata, near Sukegawa, but older in age (?). Pliocene and Miocene of Europe.

Living.—Central Japan. Northern Pacific. Atlantic.

R CB 22364

96. *Pereudesia grayi*, Davidson var. *transversa*, DAVIDSON.

Pereudesia grayi var. *transversa*. Hayasaka, Tert. Brach. Japan, p. 16, pl. II, figs. 13, 14.

Eudesia grayi. Yokoyama, Foss. Up. Musash., p. 199, pl. XVII, fig. 11, 12.

Quite frequent.

Fossil occurrence.—Tempizan. Upper Musashino. Pliocene (?) of Yanagawa-machi, Iwaki (according to Hayasaka).

Living.—Northern, Central and Western Japan. California.

INDEX.

A.		H.	
<i>Acmaea nigro-lineata</i> Rve.	15	<i>Haliotis diversicolor</i> Rve.	15
<i>Acmaea</i> sp.	15	<i>Haliotis</i> sp.	15
<i>Arca kobeltiana</i> Pils.	28	<i>Helcioniscus pallidus</i> Gld.	15
<i>Arca setensis</i> Yok.	29		
<i>Arca subcrenata</i> Lke.	28		
B.		L.	
<i>Bathybemleix argenteonitens</i> Lke.	14	<i>Leda confusa</i> Hanl.	29
<i>Batillus cornutus</i> Gm.	14	<i>Leda ramsayi</i> Sm.	30
<i>Bela rugulata</i> Tr. var. <i>schneideri</i>	9	<i>Leda</i> sp.	30
<i>Buccinum leucostoma</i> Lke.	10	<i>Lima angulata</i> Sow.	25
<i>Buccinum pericochlion</i> Schr.	10	<i>Lima goliath</i> Sow.	25
C.		<i>Lima lima</i> L.	25
<i>Calyptaea mammilaris</i> Brod.	13	<i>Lima vulgatula</i> Yok.	25
<i>Cardita cumingiana</i> Dkr.	24	<i>Limopsis azumana</i> Yok.	29
<i>Cardium californiense</i> Desh.	22	<i>Lucina borealis</i> L.	24
<i>Cardium modestum</i> A. et R.	23	<i>Lucina spectabilis</i> Yok.	24
<i>Cardium muticum</i> Rve.	23		
<i>Cardium shinjiense</i> Yok.	23	M.	
<i>Cassis japonica</i> Rve.	11	<i>Macoma dissimilis</i> Mart.	20
<i>Cerithium baculum</i> Yok.	12	<i>Macoma praetexta</i> Mart.	20
<i>Cerithium</i> sp.	12	<i>Mactra spectabilis</i> Lke.	17
<i>Chrysodomus pericochlion</i> Schr.	10	<i>Mactra sulcataaria</i> Desh.	17
<i>Chrysodomus phoeniceus</i> Dall.	10	<i>Mactra veneriformis</i> Desh.	17
<i>Chrysodomus schrencki</i> Yok.	10	<i>Mercenaria stimpsoni</i> Gld.	21
<i>Clementia speciosa</i> Yok.	21	<i>Meretrix iizukai</i> Yok.	20
<i>Conus sieboldi</i> Rve.	8	<i>Meretrix lamarekii</i> Ag.	21
<i>Conus</i> sp.	8	<i>Modiola barbata</i> L.	25
<i>Crepidula aculeata</i> Gm.	13	<i>Modiola</i> sp.	25
<i>Crepidula grandis</i> Mid.	13	<i>Mya arenaria</i> L.	16
<i>Crepidula</i> sp.	13	<i>Mya japonica</i> Jay.	16
<i>Cultellus izumoensis</i> Yok.	18	<i>Mytilus grayanus</i> Schr.	25
<i>Cyclina chinensis</i> Chem.	20	<i>Mytilus hirsutus</i> Lam.	25
D.		<i>Mytilus</i> sp.	25
<i>Dentalium complexum</i> Dall.	16		
<i>Dentalium weinkaufii</i> Dkr.	16	N.	
<i>Diplodonta semiaspera</i> Phil.	23	<i>Natica janthostoma</i> Desh.	13
<i>Dolium luteostomum</i> Küst.	11	<i>Nucula insignis</i> Ad.	30
<i>Dosinia troscheli</i> Lke.	20	<i>Nucula mirabilis</i> Ad. et Rve.	30
E.			
<i>Eudesia grayi</i> Dav.	32	O.	
G.		<i>Olivella fortunei</i> Ad.	9
<i>Galeodea japonica</i> Yok.	11	<i>Ostrea gigas</i> Thunb.	28
P.			
		<i>Panope generosa</i> Gld.	16
		<i>Parallelodon obliquatus</i> Yok.	29
		<i>Pecten laetus</i> Gld.	26

Pecten kimurai Yok.	27	
Pecten swiftii Bern.	27	
Pecten tokyoensis Tok.	27	
Pecten vesiculosus Dkr.	27	
Pecten yessoensis Jay.	27	
Pectunculus vestitus Dkr.	29	
Pereudesia grayi Dav.	32	
Phacoides borealis L.	24	
Pinna japonica Harl.	28	
Pleurotoma kamakurana Pils.	8	
Pleurotoma mediocarinata Yok.	8	
Pleurotoma sp.	8	
Polinices (Neverita) ampla Rve.	14	
Friene oregonensis Redf.	11	
S.		
Saxidomus purpuratus Low.	21	
Scapharca subcrenata Lke.	28	
Sigaretus (Runaticina) papilla Gm.	14	
Siliqua pulchella Dkr.	18	
Solarium perspectivum L.	13	
Solarium sp.	13	
Solemya tokunagai Yok.	31	
Solen gouldii Conr.	18	
Solen gracilis Gld.	18	
Solen grandis Dkr.	18	
Solen krusensternii Schr.	18	
Soletellina violacea Desh.	18	
Spisula grayana Schr.	17	
Stomatella lyrata Pils.	15	
Stomatella japonica Ad.	15	
T.		
Tapes undulatus Born.	22	
Tellina besshoensis Yok.	19	
Tellina izurensis Yok.	19	
Tellina optiva Yok.	19	
Terebra lisckeana Dkr.	8	
Terebra sp.	8	
Terebratulina caputserpentis L.	31	
Thracia pubescens Pult.	25	
Thyasira bisecta Conr.	24	
Tritonium pericochlion Schr.	10	
Trochus argenteo-nitens Lke.	14	
Turbo cornutus Gm.	14	
Turbo sp.	14	
Turritella nipponica Yok.	13	
V.		
Venericardia ferruginea Ad.	24	
Venericardia sp.	24	
Venus jedoensis Lke.	22	
Venus rimosa Phil.	22	
Venus stimpsoni Gld.	21	
Venus toreuma Gld.	22	
Venus undulatus Born.	22	
Vermetus tricarinatus Yok.	12	
Vermetus sp.	12	
Voluta megaspira Sow.	9	
Volutarpa perryi Schr.	10	
Y.		
Yoldia lisckei Sm.	30	

Errata.

P. 4, line 19 and P. 7, line 12. Read Lima goliath Sow.

M. YOKOYAMA:

Molluscan Remains from the Uppermost Part of the Jô-Ban Coal-Field.

PLATE I.

Plate I.

- Fig. 1. *Chrysodomus phoeniceus* Dall. A large specimen from Tsurushihama, Sukegawa. p. 10.
- Fig. 2. *Chrysodomus pericochlion* Schr. Cast. Tsurushihama, Sukegawa. p. 10.
- Fig. 3. *Bathybembix argenteo-nitens* Lke. Cast. Tsurushihama, Sukegawa. p. 14.
- Fig. 4. *Helcioniscus pallidus* Gld. A large cast from Tsurushihama, Sukegawa. p. 15.
- Fig. 5. *Cassis japonica* Rve. Cast. Tsurushihama, Sukegawa. p. 11.
- Fig. 6. *Clementia speciosa* Yok. Railway Cutting, Nakoso. p. 21.
- Figs. 7, 8. *Mactra spectabilis* Lke. 7. From Izura. 8. Cast from Tsuchibashi, Kadōno. p. 17.
- Fig. 9. *Pleurotoma* sp. Izura. p. 8.
- Fig. 10. *Galeodia (Scionsia) japonica* Yok. Tozenji, Yunami. p. 11.
- Fig. 11. *Thracia pubescens* Pult. Gomasawa, Taira. p. 25.





Plate II.

- Fig. 1. *Pecten swiftii* Bern. Right valve. Tsurushihama, Sukegawa. p. 27.
- Fig. 2. *Yoldia lischkei* Sm. Cast. Tsurushihama, Sukegawa. p. 30.
- Fig. 3. *Stomatella japonica* Adams. Cast. Tsurushihama, Sukegawa. p. 15.
- Fig. 4. *Pecten kimurai* Yok. Right valve. Izura. p. 27.
- Fig. 5. *Solen gouldii* Conr. Tozenji, Yunami. p. 18.
- Fig. 6. *Cerithium baculum* Yok. Yamoto, Kadōno. p. 12.
- Fig. 7. *Cerithium* sp. Yamoto. p. 12.
- Fig. 8. *Turritella nipponica* Yok. Izura. p. 13.
- Fig. 9. *Dentalium complexum* Dall. 2/1. Tempizan. p. 16.
- Fig. 10. *Mytilus grayanus* Dkr. Cast. Tempizan. p. 25.
- Fig. 11. *Spisula grayana* Schr. Cast. Tsurushihama, Sukegawa. p. 17.
- Fig. 12. *Tellina izurensis* Yok. Cast of a left valve. Izura. p. 19.



M. Yokoyama:—Molluscan Remains from the Jô-Ban Coal-Field.

Plate III.

- Fig. 1. *Lima goliath* Sow. Right valve. Kamamaye, Kadōno. p: 25.
Figs. 2, 3. *Meretrix iizukai* Yok. Fig. 2 is a specimen with the anterior end
somewhat more sharply rounded than Fig. 3. Izura. p. 20.
Fig. 4. *Lima goliath* Sow. Cast. Tsurushihama, Suwegawa. p. 25.




Plate IV.

Pecten kimurai, Yok.

P. 27.

- Fig. 1. Left valve. Izura.
- Fig. 2. Left valve seen from the inner side. Izura.
- Fig. 3. Right valve. Showing the anterior ear. Izura.
- Fig. 4. Right valve (water-worn). Tamaye, Kobisa.
- Fig. 5. Right valve. of the same individual as Fig. 1. Izura.
- Fig. 6. Right valve. Izura.




Plate V.

- Fig. 1. *Tapes undulatus* Born. Cast. Hatsuzaki in Sukegawa. p. 22.
Figs. 2, 3. *Cultellus izumoensis* Yok. 2. Nakoso Coast. 3. Iida, Kashima. p. 18.
Fig. 4. *Venericardia ferruginea* Ad. Izura. p. 24.
Figs. 5-8. *Lucina (Phacoides) borealis* L. 5. A large cast from Tsurushi-hama. 6-8. Izura. p. 24.
Fig. 9. ~~Pecten~~ *dissimilis* Mart. Kokozura. p. 20.

Macoma.





Plate VI.

- Figs. 1, 2, 3. *Solemya tokunagai* Yok. Fig. 1. A specimen from Tsuchibashi in Kadōno, showing the periostracum produced into long processes. Fig. 2. From Shiogu in Tatsuta. Fig. 3. Showing the posterior end. Enamura Quarry. p. 31.
- Fig. 4. *Mya arenaria* L. var. *japonica* Jay. Tozenji in Yunami. p. 16.
- Fig. 5. *Thyasira bisecta* Conr. Taira Park. p. 24.
- Fig. 6. *Panope generosa* Gld. Iida in Kashima. p. 16.
- Fig. 7. *Piuna japonica* Hanl. Iida in Kashima. p. 28.



Vol. XLV., Art. 5, published March 21st,
1925.

Price in Tokyo, Yen 1.75.

This Journal is on sale at
MARUZEN CO., LTD., Tori Sanchome, Nihonbashi, Tokyo.
R. FRIEDLÄNDER u. SOHN, Carlstr. II, Berlin, NW. 6.

大正十四年三月十七日印刷
大正十四年三月二十一日發行

編纂兼發行者 東京帝國大學

印 刷 者 星 野 錫

東京市日本橋區兜町二番地

印 刷 所 東京印刷株式會社

東京市日本橋區兜町二番地

賣 摆 所 丸 善 株 式 會 社

東京市日本橋區通三丁目十四番地

NOTICE

Vol. XLIII. (Concluded):

- Art. 1. K. YENDO :—A Monograph of the Genus *Alaria*. With 19 plates. Publ. May 30th, 1919.
- Art. 2. T. MATSUHIMA :—Untersuchungen über die Wasseraufnahme bei abgeschnittenen Zweigen. Mit 2 Textfiguren. Publ. July 30th, 1919.
- Art. 3. Y. YOSHII :—Oekologische Studien über Vegetation der Ota Dünen. Mit 2 Tafeln und 8 Textfiguren. Publ. July 30th, 1919.
- Art. 4. Y. EMOTO :—Über die relative Wirksamkeit von Kreuz- und Selbstbefruchtung bei einigen Pflanzen. Mit 2 Tafeln und 6 Textfiguren. Publ. March 15th, 1920.
- Art. 5. T. KATŌ :—A Contribution to the Knowledge of the Cassiterite Veins of Pneumato-Hydato-genetic or Hydrothermal Origin. A Study of the Copper-Tin Veins of the Akenobō District in the Province of Tajima, Japan. With 7 plates and 11 text figures. Publ. May 10th, 1920.
- Art. 6. S. TSUBOI :—Volcano Ōshima, Idzu. With 6 plates and 42 text figures. Publ. May 10th, 1920.
- Art. 7. M. TAHARA :—Cytologische Studien an einigen Kompositen. Mit 4 Tafeln und 15 Textfiguren. Publ. March 30th, 1921.
- Art. 8. Y. KUDŌ :—Enumeratio Labiatarum speciarum varietatum formarumque in Insulis Kurilensibus et Insula Yezoensi sponte nascentium. Publ. March 30th, 1921.

Vol. XLIV. (Concluded):

- Art. 1. M. YOKOYAMA :—ossils from the Upper Musashino of Kazusa and Shimosa. With 17 plates. Publ. July 7th, 1922.
- Art. 2. T. KABURAKI :—On some Japanese Freshwater Tricladida, with a Note on the Parallelism in their Distribution in Europe and Japan. With 1 plate and 14 text figures. Publ. June 23rd, 1922.
- Art. 3. T. KABURAKI :—On some Japanese Tricladida Maricola, with a Note on the Classification of the Group. With 1 plate and 6 text figures. Publ. September 7th, 1922.
- Art. 4. T. KABURAKI :—On the Terrestrial Planarians from Japanese Territories. With 1 plate and 23 text figures. Publ. September 30th, 1922.
- Art. 5. T. TAKAGI :—Ueber das Reciprocitätsgesetz in einem beliebigen algebraischen Zahlkörper. Publ. September 16th, 1922.
- Art. 6. T. TERADA :—On some Remarkable Relations between the Yearly Variations of Terrestrial Phenomena and Solar Activities. With 17 text figures. Publ. January 24th, 1923.
- Art. 7. M. YOKOYAMA :—On some Fossil Shells from the Island of Saishū in the Strait of Tsushima. With 1 plate. Publ. January 24th, 1923.
- Art. 8. E. STECHOW :—Die Hydroidenfauna der Japanischen Region. Publ. February 26th, 1923.

Vol. XLV.:

- Art. 1. M. YOKOYAMA :—Mollusca from the Coral-Bed of Awa. With 5 plates. Publ. October 3rd, 1924.
- Art. 2. M. YOKOYAMA :—Tertiary Mollusca from Dainichi in Tōtōmi. With 2 plates. Publ. March 31st, 1923.
- Art. 3. M. YOKOYAMA :—Molluscan Remains from the Lowest Part of the Jō-Ban Coal Field. With 5 plates. Publ. December 30th, 1924.
- Art. 4. Y. OZAWA :—On the Classification of Fusulinidae. With 4 plates and 3 text figures. Publ. February 22nd, 1925.
- Art. 5. M. YOKOYAMA :—Molluscan Remains from the Uppermost Part of the Jō-Ban Coal-Field. With 6 plates. Publ. March 21st 1925.