

139

June 25th, 1925.

Vol. XLV., Art. 7.

東京帝國大學  
理學部紀要

1925b

第四拾五冊 第七編

18

JOURNAL  
OF THE  
COLLEGE OF SCIENCE,  
IMPERIAL UNIVERSITY OF TOKYO.

22367  
22539

MATAJIRO YOKOYAMA.

Molluscan Remains from the Middle Part of the  
Jô-Ban Coal-Field.

27092  
27095

TOKYO.

PUBLISHED BY THE UNIVERSITY.

TAISHO XIV.

欠

Publishing Committee.

Prof. S. Goto, *Rigakuhakushi*, Director of the College (*ex officio*).

Prof. K. Matsubara, *Rigakuhakushi*.

Prof. K. Fujii, *Rigakuhakushi*.

Prof. N. Yamasaki, *Rigakuhakushi*.

Prof. N. Yatsu, *Ph. D.*, *Rigakuhakushi*.



All communications relating to this Journal should be addressed to the  
Director of the College of Science.

## Molluscan Remains from the Middle Part of the Jô-Ban Coal-Field.

By

Matajiro YOKOYAMA *Rigakuhakushi.*

---

*With 3 Plates.*

---

In the two papers<sup>1)</sup> recently published on the molluscan remains of the Jô-Ban coal-field, I described those from the lowest beds as well as from the uppermost. What now remains are those of the beds between, which are treated in the following pages.

Unfortunately, the fossils sent me by Dr. Tokunaga as derived from these beds are not only small in number, but very imperfectly preserved, making their determination extremely difficult. Moreover, while the already known intermediate beds containing fossils are four in number—in descending order, the Misawa, the Kamenoo, the Mizunoya and the Goyasu<sup>2)</sup>—what I have obtained are only the specimens from two of them, the *Kamenoo* and the *Mizunoya*. Besides, however, there are also fossils from beds whose exact geological position is at present unknown, but which are decidedly younger than the above and much better preserved.

### 1. Fossils from the Mizunoya and Kamenoo-Beds.

The fossil-bearing rock is a shale grey to yellowish in colour and often sandy or tufaceous. The fossils are mostly flat-pressed or distorted.

The species from the *Mizunoya-Beds* which I could distinguish are the following :

1. *Turritella nipponica* Yok.
2. *Turritella* sp.
3. *Tellina besshoensis* Yok.
4. *Tellina* sp.

---

1) Jour. Sci. Coll., Vol. XLV, Art. 3 and Art. 5.

2) The fifth beds, the Shirasaka, which are below the Goyasu, do not contain fossils.

45. *Lucina* (*Phacoides*) *borealis* L.
6. *Venericardia ferruginea* Ad.
7. *Pectunculus* sp.
8. *Yoldia sagittaria* n. sp.
9. *Nucula insignis* Ad.

Leaving out 3 which are not specifically determined, there remain only 6, of which *Tellina besshoensis* is *Miocene* and *Pliocene*, *Lucina borealis*, *Venericardia ferruginea* and *Nucula insignis* are *Miocene*, *Pliocene*, and *recent*. *Turritella nipponica* is *Pliocene*, while *Yoldia sagittaria* is entirely new. Therefore, if we are left with these few forms to judge their age, we may say they are either *Miocene* or *Pliocene*, though a little more weight seems to fall on the latter determination. Such being the case, it may be advisable to see how things are in the beds above, the *Kamenoo*. Here the number of species is somewhat greater and consequently better fitted to determine their age. They are the following:

1. *Natica* sp.
2. *Macoma dissimilis* Mart.
3. *Lucina* (*Phacoides*) *borealis* L. (?)
4. *Lucina* (*Phacoides*) sp.
5. *Venericardia laxata* Yok.
6. *Venericardia pacifera* Yok. (?)
7. *Venericardia ferruginea* Ad.
8. *Venericardia orbica* n. sp.
9. *Pecten tairanus* n. sp.
10. *Pectunculus* sp.
11. *Leda pennula* n. sp.
12. *Leda inermis* n. sp.
13. *Yoldia tokunagai* n. sp.
14. *Nucula eximia* n. sp.
15. *Solemya tokunagai* Yok.

Out of 10 specifically well determined forms, 1 (*Venericardia laxata*) is *Miocene*, 1 (*Venericardia ferruginea*) *Miocene*, *Pliocene* and *recent*, 1 (*Macoma dissimilis*) *Pliocene* and *recent*, 1 (*Solemya tokunagai*) up to this time only *Pliocene*, while 6 are quite new. From this, we can not but consider the *Kamenoo-Beds* as *Miocene*, from which it naturally follows that the *Mizunoya-Beds* which underlie the *Kamenoo* must also be

*Miocene.*

## 2. Fossils from Beds whose Geological Position is Still Uncertain.

In and near the sea-side town of Minato in Hitachi and also in the regions of Shimo-Ogawa and Tateishi in Iwaki, Dr. Tokunaga found fossiliferous beds whose exact geological position he is unable to point out, that is to say, to which of the several beds, already distinguished near Yumoto they correspond. The rock is mostly sandstone, tolerably hard, dark-grey to light grey and fine-grained, though sometimes it may become coarse, breccia-like and greenish as at Hiraiso, or include pebbles and become a real conglomerate as near the Matsugane-Mine. Anyhow, petrographically, the beds show some peculiar character, foreign to the already known ones.

The fossil localities in the Minato Region are given as follows: Minato, Ushikubo, Kaimon Bridge, Isozaki and Hiraiso.<sup>1)</sup>

In the Shimo-Ogawa Region,<sup>2)</sup> the beds occur at Shibahara, Ochiai, Tateshitazawa and Matsugane Mine. The fossil-bearing rock of the first two places is mostly similar to that of Minato, that is to say, a hard dark-grey sandstone, though from Shibahara there are a few pieces of yellowish shale containing fossils (only two species, a *Tellina* and *Cardium shinjiense*). The rock of Tateshitazawa is a soft grey shale with only two species of fossils, *Cardium shinjiense* and *Phacoides borealis*, while that of the Matsugane Mine is a coarse conglomerate with a whitish sandy cement, on which account the fossils are very badly preserved.

In the Tateishi Region,<sup>3)</sup> the fossil locality is only one; that is a place between the villages of Tateishi and Shiogu. The rock is a light-grey fine-grained sandstone in which rounded pebbles of tufaceous rock, are scattered.

The number of fossil species found in all these places amounts to 42, as seen in the following table:

- 
- 1) 常陸那珂郡湊、牛久保、海門橋、磯崎、平磯。
  - 2) 磐城石城郡下小川、柴原、落合、立下澤、立平久保松金嶺山。
  - 3) 同双葉郡龍田村立石、鹽具ノ間。

	Minato-Region	Tateishi-Region	Shimo-Ogawa-Region	Occurrence in Japan.
1. Ringicula sp.	+			
2. Pleurotoma sp.	+			
3. Fusus perplexus Ad.	+			Pliocene—Recent.
4. Chrysodomus despectus L.			+	Pliocene—Recent.
5. Nassa (Hima) japonica Ad	+			Pliocene—Recent.
6. Triton (Priene) oregonensis Redf.	+	+		Pliocene—Recent.
7. Cassis japonica Rve.	+			Pliocene—Recent.
8. Turritella nipponica Yok.	+		+	Pliocene.
9. Calyptraea mammillaris Brod.	+			Miocene—Recent.
10. Crepidula aculeata Gm.	+			Recent.
11. Crepidula convexa n. sp.	+			
12. Natica janthostoma Desh.		+	+	Miocene—Recent.
13. Polinices (Neverita) ampla Phil.	+			Pliocene—Recent.
14. Bembix sp.		+		
15. Fissuridea humilis n. sp.	+			
16. Chlorostoma sp.			+	
17. Helcioniscus pallidus Gld.			+	Pliocene—Recent.
18. Dentalium sp.	+			
19. Panope generosa Gld.	+	+		Pliocene—Recent.
20. Mactra sulcataria Desh.	+			Pliocene—Recent.
21. Spisula grayana Schr.	+			Pliocene—Recent.
22. Racta elliptica Yok.	+			Pliocene (or Pleistocene).
23. Tresus sp.		+		
24. Tellina sp.			+	
25. Macoma inquinata Desh.	+			Miocene—Recent.
26. Dosinia troscheli Lke.	+	+	+	Pliocene—Recent.
27. Dosinia angulosa Phil.	+		+	Recent.
28. Venus (Mercenaria) stimpsoni Gldi.		+		Pliocene—Recent.
29. Tapes variegatus Hanl.	+			Pliocene (or Pleistocene)—Recent.
30. Cardium shinjiense Yok.	+		+	Miocene—Pliocene.
31. Diplodonta usta Gld.	+		+	Pliocene (or Pleistocene)—Recent.
32. Lucina (Phacoides) borealis L.	+	+	+	Miocene—Recent.
33. Venericardia ferruginea Ad.	+			Miocene—Recent.
34. Cardita cumingiana Dkr.	+			Pliocene (or Pleistocene)—Recent.
35. Lima goliath Sow.		+		Pliocene—Recent.
36. Pecten laetus Gld.	+			Pliocene—Recent.
37. Pecten iwakianus n. sp.		+		
38. Arca sp.	+			
39. Pectunculus vestitus Dker.	+			Miocene—Recent.
40. Parallelodon obliquatus Yok.	+			Miocene—Recent.
41. Yoldia sagittaria Yok.		+		Miocene (Mizunoya).
42. Nucula mirabilis Ad. et Rve.			+	Pliocene—Recent.

The region which yielded the greatest number of species is Minato with 30. The Shimo-Ogawa and Tateishi-Regions gave respectively only 12 and 11. There are no noteworthy peculiarities in these little faunas except the great frequency of *Turritella nipponica* and *Crepidula aculeata* in the Minato-Region and of *Phacoides borealis* and *Nucula mirabilis* in the Shimo-Ogawa.

The proportion of extinct forms against living is in all the three regions nearly equal, that is to say about 20%. If we try to judge the geological age of the beds in which these fossils were found from this percentage as well as from other considerations given in the last column of the above table, we can not but say that it is *Pliocene*, as the greatest weight seems to fall on that age. In fact, palaeontologically speaking, the whole fauna can hardly be distinguished from that of the Shirado-Beds which I took for *Upper Pliocene*. Nevertheless, there is one reason which forces me to assume that the beds in question, in spite of their belonging to the same age, are lower in position than the Shirado. And that reason rests on a careful observation of Dr. Tokunaga in the Minato-Region. Here the beds are said to be markedly inclined, while the layers above which he takes for the Shirado are nearly horizontal, thus showing a distinct line of unconformability between the two.

From this it may perhaps be questioned if these inclined beds of Minato are not the equivalent of the Misawa of the Yumoto-Region which conformably overlies the Kamenoo, but unconformably underlies the Shirado. To this question it is at present not possible to answer, as the fossil content of the Misawa is still quite unknown, although petrographically there is a difference. Anyhow, when we consider the great resemblance of the Minato fossils to those of the Shirado and their little relation to those of the Kamenoo, it is highly probable that they belong to a horizon, if not quite the same, still very close to the Shirado-Beds, which in the neighbourhood of Yumoto did not come to deposition.

It is to be mentioned here that at present it is still not quite certain whether the Shimo-Ogawa and Tateishi-beds are strictly contemporaneous with those of Minato, though petrographically viewed, they presumably are.

## Description of the Species.

### 1. Fossils from the Mizunoya and Kamenoo-Beds.

#### 1. *Turritella nipponica*, Yokoyama

*Turritella nipponica*. Yokoyama, Foss. Miura Penin., p. 71, pl. IV, figs. 16—19. Moll. Rem. Uppermost Part Jō-Ban Coal-field, p. 13, pl. II, fig. 8.

CM 22367  
CM 22368

A few examples.

Fossil occurrence.—*Mizunoya-Beds*: Fujiwara.<sup>1)</sup> Shirado-Beds. Lower Musashino.

CM 22369-1-1

CM 22370

## 2. *Turritella* sp.

Pl. I Fig. 1.

Casts of a large form whose spiral ridges are quite variable in number and size. Some have five subequal ridges, while others have the lower two very broad and transversely striated, while upper ones are comparatively very small. All the specimens are in fragments, so that it is impossible to make out their exact characters. There is a doubt, however whether they do not belong to an abnormally large form of the preceding species.

CM 22371

CM 22372

Fossil occurrence.—*Mizunoya-Beds*: Nagako in Kubota-mura.<sup>2)</sup>

## 3. *Natica* sp.

A very poorly preserved specimen. The species is not determinable.

Fossil occurrence.—*Kamenoo-Beds*: Fukuda,<sup>3)</sup> Sekimoto.

CM 22373-1-19

CM 22374

CM 22375

## 4. *Macoma dissimilis*, Martens (?)

Pl. I, Fig. 19.

A few casts closely resembling in form to the above named species already found in the Shirado-beds (Yokoyama, Moll. Rem. Uppermost Part Jō-Ban Coalf., p. 20, pl. v. fig. 9).

Fossil occurrence.—*Kamenoo-Beds*: Yunami<sup>4)</sup>; between Yunami and Onsenba. Shirado-Beds. Musashinos.

CM 22376

CM 22377

CM 22378

## 5. *Tellina besshoensis*, Yokoyama.

*Tellina besshoensis*. Yokoyama, Moll. Rem. Lowest Part Jō-Ban Coalfield, p. 14, pl. II, figs. 1—5. Moll. Rem. Uppermost Part, p. 19.

Several undoubted specimens.

Fossil occurrence.—*Mizunoya-Beds*: Fujiwara; Yuzen,<sup>5)</sup> Yunami. Asagai-Beds. Shirado-Beds.

CM 22379

## 6. *Tellina* sp.

An indeterminable form.

Fossil occurrence.—*Mizunoya-Beds*: east of Tebaizaka in Yumoto.<sup>6)</sup>

1) 磐城石城郡磐崎村藤原 (大日本炭礦會社事務所下)

2) 同郡窪田村長子

3) 常陸多賀郡關本村福田

4) 常陸多賀郡關本村湯網

5) 同湯善

6) 湯本手這坂ノ東



7. *Lucina* (*Phacoides*) *borealis*, Linné.

Pl. I. Fig. 2 (?)

*Lucina borealis*. Yokoyama, Foss. Miura Penin., p. 133, pl. X, fig. 7. Moll. Rem. Uppermost Part Jō-Ban Coalfield, p. 24, pl. V, figs. 5-8.

A right valve from the Mizunoya-Beds tolerably large and measuring 50 millim. in length and 49 millim. in height belongs undoubtedly to this species. There are also specimens from the Kamenoo-Beds like the one here figured which, though imperfect, seem to belong to the same species.

Fossil occurrence.—*Mizunoya-Beds*: Fujiwara. *Kamenoo-Beds* doubtful specimens), Fukuda and Yunami.

Living.—Central Japan.

8. *Lucina* (*Phacoides*) sp.

Pl. I. Figs. 3, 4, 5, 18.

There are several flatly pressed impressions of a *Lucina* which are apparently different from the preceding species in being generally longer and more markedly subtruncate behind. The sculpture is similar, showing distant, raised, concentric lamellae with fine concentric lines between. The proportion of length to height is variable, but the shell is invariably longer than high. Whether these specimens belong to a distinct species, or are only the deformations of the preceding, I am now not in a position to decide.

Fossil occurrence.—*Kamenoo-Beds*: Ebata (Yamada)<sup>1)</sup>; Ishibatake (Nishiki)<sup>2)</sup>; Ishimoriyama<sup>3)</sup>; Tabasaka (Yumoto)<sup>4)</sup>; Tsūchibashi (Kadono)<sup>5)</sup>; Miyanosaku (Shimoyata near Yumoto)<sup>6)</sup>; Ōdate, Taïra.<sup>7)</sup>

9. *Venericardia laxata*, Yokoyama.

Pl. I. Figs. 11, 12 (?)

*Venericardia laxata*. Yokoyama, Moll. Rem. Lowest Part Jō-Ban Coal-Field, p. 19, pl. III, figs. 16-18.

A rather small, more or less transversely roundly elliptical shell with flat radiating ribs which number a little over twenty is, I believe, the same species as that described from the Asagai-beds, though somewhat shorter in form.

Fossil occurrence.—*Kamenoo-Beds*: Shimo-Yunagaya (Yumoto).<sup>8)</sup> Doubtful specimens are found at Ebata (Yamada), Tanamisaku (Shimofunao),<sup>9)</sup> and Hirakata coast.<sup>10)</sup>

- 1) 磐城石城郡山田村江畑 2) 同錦村石畑 3) 石森山一名磐城富士 (平町ノ北、平窪村)  
4) 田場坂 5) 上遠野土橋 6) 下矢田宮ノ作 7) 平町大館 8) 湯本下湯長谷  
9) 下船尾蛇井作 (磐崎村) 10) 平沼海岸

CM 22380  
CM 22381-1-2  
CM 22382  
CM 22383

CM 22384  
CM 22385  
CM 22386-1-3  
CM 22387  
CM 22388-1-18

CM 22389-1-5  
CM 22390  
CM 22391  
CM 22392-1-4

CM 22392  
CM 22393-1-11  
CM 22394  
CM 22395-1-12  
CM 22396  
CM 22397  
CM 22398

CM 22 399-1-13

10. *Venericardia pacifera*, Yokoyama (?)

Pl. I. Fig. 13.

A young individual with about fifteen ribs, distant and flat, seems to belong to the above-named species (Moll. Rem. Lowest Part., p. 18, pl. IV, figs. 1, 2.)

Fossil occurrence.—*Kamenoo-Beds*: Daikatsu Bore-hole at Yokone-sawa (Uchigō).<sup>1)</sup>

CM 22 400

CM 22 401-1-6

CM 22 402-1-7

CM 22 403

*Venericardia ferruginea*, Adams.

Pl. I. Figs. 6, 7.

*Venericardia ferruginea*. Yokoyama, Moll. Rem. Upper most Part, p. 24, pl. V, fig. 4.

*Venericardia* sp. Yokoyama, Moll. Rem. Lowest Part, p. 19, pl. III, figs. 8, 9.

Poorly preserved specimens.

Fossil occurrence.—*Mizunoya-Beds*: Fujiwara. *Kamenoo-Beds*: Tsuchibashi (Kadono).<sup>2)</sup> *Shirado-Beds*. Musashino Formation.

Living.—Northern Japan.

CM 22 404-1-10

12. *Venericardia orbica*, n. sp.

Pl. I. Fig. 10.

An ill-preserved specimen which, however, is characteristic enough to be created into a new species.

Shell nearly circular, almost as high as long, ornamented with nearly thirty flattish radiating ribs separated by interspaces of nearly equal breadth. Growth-lines distinct, fine, numerous and crossing the ribs. Length 34 millim. Height 32 millim. The thickness is not measurable, as the only specimen we possess is flatly pressed on the stone.

This shell has a striking resemblance in form to *Venericardia imbricata* Lam. of the Paris Basin as reproduced in Zittel's Text-book of Paleontology. p. 475 (fig. 764).

Fossil occurrence.—*Kamenoo-Beds*: Ishibatake (Nishiki-mura).

CM 22 405-1-8

CM 22 406-1-9

13. *Pecten tairanus*, n. sp.

Pl. I. Figs. 8, 9.

A small, neat, flat shell circular in outline, slightly longer than high. The surface is smooth, only coarse concentric corrugations being present. The ears of the left valve, though somewhat unequal in shape, are triangular, while the anterior one of the right valve is long and narrow with a few coarse radiating ribs and a deep byssal notch below. The left valve

1) 内郷村横根澤大勝試錐孔 2) 上遠野土橋 3) 平町久保町

(fig. 8) which is nearly perfect is 13 millim. long and 11 millim. high.

Fossil occurrence.—*Kamenoo-Beds*: Kubo-Street in the town of Taira.

CM 22407  
CM 22408-1-17  
CM 22409

#### 14. *Pectunculus* sp.

Pl. I. Fig. 17.

Imperfect specimens, much like those of *Pectunculus vestitus* Dkr. (Yokoyama, Upper Musashino, pl. XVI, figs. 1, 2).

Fossil occurrence.—*Mizunoya-Beds*: East of Tebaizaka, Yumoto.  
*Kamenoo-Beds*: Mayanosaku, Shimoyata

CM 22410-2-9  
CM 22411  
CM 22412  
CM 22413-2-8  
CM 22414

#### 15. *Leda pennula*, n. sp.

Pl. II. Figs. 7, 8, 9.

Shell small, thin, compressed, transversely elongated, very inequilateral, rounded in front, longly rostrate, narrowed, and mostly truncate behind. The length is about double the height. Postero-dorsal border nearly straight, inclining to concave; ventral almost straight in the posterior half. Area narrow, distinct. Surface with prominent concentric lirae, suddenly turned upward at the areal border with a broad angle and continuing into the area. A few examples, one of which measures 19 millim. long and 9 millim. high.

This species looks like *Leda ramsayi* Sm. (Foss. Miura Penin., p. 176, pl. XIX, fig. 3), but the shell is longer.

Fossil occurrence.—*Kamenoo-Beds*. Daikatsu bore-hole, Yokone-sawa; Yotsusawa (near Nakoso Station), Kubota<sup>1)</sup>; Takinosaku (Kami-Arakawa, Iino-mura).<sup>2)</sup> A doubtful specimen occurs also at Saku (Iino).<sup>3)</sup>

CM 22417-2-4  
CM 22418  
CM 22419-2-6  
CM 22420  
CM 22421-2-3  
CM 22422-2-5

#### 16. *Leda inermis*, n. sp.

Pl. II. Figs. 1-6.

Shell comparatively large, thin, flatly convex, transversely oblong, with the length usually a little over one and a half times the height; inequilateral with anterior side somewhat shorter than posterior; rounded in front, rounded or subtruncate behind, broadly arched at ventre; antero- and postero-dorsal borders arched, the former more sharply than the latter. Surface smooth. Teeth about 20 on the posterior side, about half as many on the anterior.

CM 22424-2-1  
CM 22425-2-2  
CM 22426  
CM 22427

1) 窪田四澤 (石城郡) 2) 飯野村上荒川瀧ノ作 3) 飯野村作

The largest specimen measures a little over 40 millim. in length. The thickness of the shell can not be well measured as it is more or less flattened by pressure, but it seems to be about one-sixth the length. Rather frequent.

Fossil occurrence.—*Kamenoo-Beds*: Tamatsuyu (Izumi)<sup>1)</sup>; Hirakata coast; Inouye (Uyeda)<sup>2)</sup>; Osaku near Nakoso Railway Station<sup>3)</sup>; between Yunami and Onsenba.

CM 22428-2-11

CM 22429

CM 22430-2-10

CM 22431

17. *Yoldia sagittaria*, n. sp.

Pl. II. Figs. 10, 11.

Shell thin, strongly compressed, lanceolate, more than twice as long as high, subequilateral, anteriorly rounded, posteriorly narrowed, rostrate and pointed (?) Surface with distant concentric impressed lines. Teeth about 25 on the anterior side, a little less on the posterior. Rare.

Two specimens. The one measures 22 millim. high and 45 (?) millim. long. The other is 27 millim. high and more than 50 millim. long.

Fossil occurrence.—*Mizunoya-Beds*: Between Kosai and Aburahira (Sekimoto)<sup>4)</sup>; Nagako, Kubota.

CM 22432

CM 22433

CM 22434-2-17

CM 22435

CM 22436-2-15

CM 22437-2-18

CM 22438

CM 22439-2-73

CM 22440

CM 22441-2-16

CM 22442

CM 22443-2-12

CM 22444

CM 22445

CM 22446

CM 22447

18. *Yoldia tokunagai*, n. sp.

Pl. II. Figs. 12-18.

Shell medium-sized, thin, compressed, transversely suboblong, nearly twice as long as high, inequilateral with the anterior side almost double the length of the posterior, rounded in front, subrostrate and rounded behind: antero-dorsal border nearly straight, postero-dorsal slightly excavated, ventral broadly arched and obliquely ascending behind. Surface with fine divaricating lines. Anterior teeth a little over 20 in number, posterior a little less and coarser.

This species is somewhat like *Yoldia laudabilis* Yok. (Moll. Rem. Lowest Part., p. 22. pl. IV. figs. 11, 12) of the Asagai-Beds in outline, though the surface-sculpture is quite different.

Fossil occurrence.—*Kamenoo-Beds*: Ishimoriyama; Kami-Kabeya<sup>5)</sup>; Saku; Takinosaku; Miyanosaku; Honya (Izumi)<sup>6)</sup>; Shimoyunagaya (Yumoto); between Yunami and Onsenba. A doubtful specimen occurs also at Yotsusawa, Kubota.

CM 22448

CM 27092

(R)

19. *Nucula insignis*, Adams.

*Nucula insignis*. Yokoyama, Moll. Rem. Lowest Part Jō-Ban Coal-Field, p. 21. Moll. Rem. Uppermost Part, p. 30.

1) 泉村玉露 (石城郡)

2) 植田井上

3) 大作

4) 關本村小才油平ノ間 (多賀郡)

5) 上神谷 (神谷村)

6) 泉村本谷

An imperfect specimen.

Fossil occurrence.—*Mizunoya-Beds*: Yuzen, Yunami<sup>1)</sup>; Asagai-Beds.

Shirado-Beds. Musashinos.

Living.—Northern Japan.

CM 22449-1-15

CM 22450

CM 22451

CM 22452-1-16

CM 22453

CM 22454-1-14

## 20. *Nucula eximia*, n. sp.

Pl. I. Figs. 14, 15, 16.

CM 22455

Shell rather large, trigonal, somewhat longer than high, a little inequilateral, anterior side longer than posterior, rounded at both ends, although sharply in front and quite broadly behind; antero-dorsal border nearly straight, sloping, postero-dorsal but slightly arched and also descending. Surface with neat divaricating riblets. The proportion of length to height is 10 to about 8.2. The largest example measures some 40 millim. in height.

This species is quite characteristic by its large size, trigonal shape and divaricating sculpture.

Mostly found as casts and flatly pressed, so that it is not possible to know the thickness.

Fossil occurrence.—*Kamenoo-Beds*: Hayashizaku (Iino); Takinosaku (Iino); Saku (Iino); Kamenoo-sawa (Mizunoya); north of Sekiba (hill-top).

CM 22456a-1-20

CM 22456b-1-21

## 21. *Solemya tokunagai*, Yokoyama.

Pl. I. Figs. 20, 21.

*Solemya tokunagai*. Yokoyama, Moll. Rem. Upper. Part, p. 31, pl. VI, figs. 1-3.

Imperfect specimens. Yet they are easily recognized by the peculiar processes of the periostracum.

Fossil occurrence.—*Kamenoo-Beds*: west of Fukuda. Shirado-Beds.

## II. Fossils from Uncertain Beds.

CM 22457

### 1. *Ringicula* sp.

A few ill-preserved specimens resembling *Ringicula musashinoensis* Yok. (Foss. Miura Penin., p. 30, pl. I., figs. 3, 8), but somewhat more slender.

Fossil occurrence.—*Minato Region*: Ushikubo.

1) 湯網湯善

CM 22458

2. *Pleurotoma* sp.

Two mutilated specimens, fusiform in shape and having three spiral ridges on the whorls except the last which has eight.

Fossil occurrence.—*Minato Region*: Ushikubo.

CM 22459

3. *Fusus perplexus*, A. Adams.

*Fusus perplexus*. Yokoyama, Foss. Miura Penin., p. 50, pl. II, fig. 17. Foss. Up. Musash., p. 51.

A fragmentary example.

Fossil occurrence.—*Minato Region*: Ushikubo. Musashinos.

Living.—Northern, Central and Western Japan.

CM 22460

4. *Chrysodomus despectus*, (Linné).

*Neptunea despecta*. Harmer, Plioc. Moll. Brit. Brit., part I, p. 160.

*Chrysodomus phoeniceus*. Yokoyama, Foss. Miura Penin., p. 50, pl. II, figs. 8—10, Moll. Rem. Lowest Part Jō-Ban Coal., p. 9. Moll. Rem. Upper Part, p. 10, pl. I, fig. 1.

The shell which I used to call *Chrysodomus phoeniceus* Dall is, as I am now convinced, identical with *Neptunea despecta* L. of Harmer. This author distinguishes some ten varieties out of this variable species, but I must confess that it is not always possible to determine to which variety or varieties our fossil specimens belong, especially when their preservation is not perfect. Nevertheless, I may say that those from the Musashino of Koshiba (figs. 8-10, pl. II, Foss. Miura Penin.) agree with such varieties as *carinata*, *antiquata* and *intersculpta* of Harmer. A specimen figured in plate I of my "Molluscan Remains from the Uppermost Part of the Jō-Ban Coal-field" seems to resemble var. *intersculpta* most.

Fossil occurrence.—*Shimo-Ogawa Region*: Matsugane Mine. Shirado-Beds. Lower Musashino.

Living.—Circumpolar seas. Northern down to Western Japan.

CM 22461

5. *Nassa (Hima) japonica*, A. Adams.

*Nassa (Hima) japonica*. Yokoyama, Foss. Miura Penin., p. 56, pl. II, fig. 5, Foss. Up. Musash., p. 58.

Several specimens, though imperfect.

Fossil occurrence.—*Minato Region*: Ushikubo. Musashinos.

Living.—Central and Western Japan.

CM 22462

CM 22463

6. *Triton (Priene) oregonensis*, Redfield.

*Triton (Priene) oregonensis*. Yokoyama, Foss. Miura Penin., p. 64, pl. III, figs. 19, 20. Foss. Up. Musash., p. 68. Moll. Rem. Up. Part Jō-Ban Coal., p. 11.

Some fragments.

Fossil occurrence.—*Minato Region*: Isozaki. *Tateishi-Region*: Between Tateishi and Shioga. Shirado-Beds. Musashinos.

Living.—Northern Japan, Alaska, Oregon, etc.

CM 22464

#### 7. *Cassis japonica*, Reeve.

*Cassis japonica*. Yokoyama, Moll. Rem. Upper. Part Jô-Ban Coal., p. 11, pl. I, fig. 5.

A cast.

Fossil occurrence.—*Minato Region*: Hiraiso. Shirado-Beds.

Living.—Central and Western Japan.

~~CM 22465~~  
CM 22465-3-5  
CM 22466-3-5  
CM 22467  
CM 22468

#### 8. *Turritella nipponica*, Yokoyama.

Pl. III. Figs. 4, 5.

*Turritella nipponica*. Yokoyama, Foss. Miura Penin., p. 71, pl. IV, Figs. 16—19. Moll. Rem. Upper. Part., p. 13, pl. II, fig. 8.

Specimens are quite frequent and found in almost all the fossil localities.

CM 22469  
CM 22470  
CM 22471  
CM 22472  
CM 22473

When I first described this species from the Musashino Formation, the specimens obtained were rather young ones which normally had four spiral ridges, though abnormally five or even six. In the specimens which I now possess, there are many young as well as fully grown ones in which latter especially an interstitial ridge appears between the main ones. These interstitials often grow in size so as to make the whorls appear several ridged. In such cases, interstitial striae may again appear, so that the number of ridges which are of various sizes becomes many, and especially on the body-whorl.

Fossil occurrence.—*Minato Region*: Ushikubo, Minato, Kaimon Bridge and Isozaki. *Shimo-Ogawa Region*: Shibahara, Yachishita, Ochiai. Shirado-Beds. Lower Musashino.

CM 22474  
CM 22475

#### 9. *Calyptrea mammilaris*, (Broderip).

*Calyptrea mammilaris*. Yokoyama, Moll. Rem. Lowest Part., p. 11, pl. I, fig. 17. Moll. Rem. Uppermost Part., p. 13.

Rather frequent.

Fossil occurrence.—*Minato Region*: Ushikubo and Hiraiso. Asagai-Beds. Shirado-Beds.

CM 22476-3-8  
CM 22477-3-9  
CM 22478  
CM 22479

#### 10. *Crepidula aculeata*, (Gmelin).

Pl. III, Figs. 8, 9.

*Crepidula aculeata*. Pilsbry, Catalogue Marine Mollusks Japan, p. 70. Tryon, Man. Conch., VIII, pl. 39, figs. 61—65. Iwakawa, Cat. Jap. Moll. Tokyo Imp. Museum, p. 49.

Numerous examples from Hiraiso. All are more or less decorticated. Now and then, there are specimens with traces of radiating ribs which are occasionally divaricating, a character also sometimes seen in the living ones. The sculpture according to the figures of Tryon seems to be very variable.

Fossil occurrence.—*Minato Region*: Hiraiso and Ushikubo.

Living.—Central, Western and Southern Japan. Australia, India, West coast of America, Florida, etc.

CM22480-3-3

# 11. *Crepidula convexa*, n. sp.

Pl. III. Fig. 3.

A single specimen which, though to a greater part deprived of its shell, is characteristic enough to be described as a new species.

It is oval in outline and high-convex, the length being 30 millim., the breadth 18 millim. and the height 14 millim. The apex is small, prominent, narrowed and recurved. The surface, so far as can be judged from a little of the shell remaining, seems to have been smooth, being only corrugated by rude lines of growth.

The general form of this shell strongly reminds of *Crepidula unjana*. Dall (Geol. and Palaeont., Harriman Alaska Expedition, p. 119, pl. x, figs. 8, 9) from the Miocene of Alaska which has the apical part more inflated.

Fossil occurrence.—*Minato Region*: Ushikubo.

CM22481

CM22482

CM22483

# 12. *Natica janthostoma*. Deshayes.

*Natica janthostoma*, Yokoyama, Moll. Rem. Lowest Part., p. 12, pl. I, fig. 20. Moll. Rem. Upper. Part., p. 13.

Fossil occurrence.—*Shimo-Ogawa Region*: Shibahara and Ochiai. (frequent). *Tateishi-Region*: Between Tateishi and Shiogu. Asagai-Beds. Shirado-Beds. Lower and Upper Musashino.

Living.—Northern and Central Japan. Kamchatka.

CM22484

CM22485

# 13. *Polinices (Neverita) ampla*, (Philippi).

*Polinices ampla*. Yokoyama, Foss. Miura Penin., p. 78, pl. V, figs. 5, 6. Foss. Up. Musash., p. 84. Moll. Rem. Upper. Part., p. 14.

Rare.

Fossil occurrence.—*Minato Region*: Ushikubo. and Kaimon Bridge. Shirado-Beds. Lower and Upper Musashino. Pliocene of Kii and Totomi.

Living.—Northern, Central and Western Japan. Philippines.



14. *Bembix* sp.

CM 22486

A fragment of the body-whorl, much decorticated and looking like that of *Bembix crumpii* Pils. (Yokoyama, Foss. Miura Penin., pl. v. figs. 27, 28.) living in Japan and also fossil in the Musashino Formation.

Fossil occurrence.—*Tateishi Region*: Between Tateishi and Shiogu.

15. *Fissuridea humilis*, n. sp.

CM 22487-3-7

Pl. III. Fig. 7.

Shell rather thin, oblong in outline, low-conical; the apex in front of the middle; anterior slope almost straight, shorter than the slightly arched posterior one which has a blunt edge in the middle. The sculpture consists of numerous, subequal, straight, radiating riblets occasionally with a finer interstitial between; the interspaces are finely latticed by growth-lines.

A single example not quite perfect, about 22 millim. long, 13 millim. broad and 5 millim. high. The shape of the apical fissure can not be well made out, as it is somewhat broken.

This shell has a close resemblance to *Fissuridea patagonica* Orb. (Tryon, Man. Conch., XII, pl. 61, figs 21—23). now living in Patagonia. It is, however, distinguished by its shape being not narrowed in front and lower.

Fossil occurrence.—*Minato-Region*: Hiraiso

CM 22488

16. *Chlorostoma* sp.

Several casts which seem to belong to *Chlorostoma argyrostoma* Gm. (Lischke, Jap. Meeresconch., I, p. 96, pl. VII, figs. 3—5) living in our seas.

Fossil occurrence.—*Shimo-Ogawa Region*: Matsugane Mine.

17. *Helcioniscus pallidus*, (Gould).

CM 22489

*Helcioniscus pallidus*. Yokoyama, Foss. Miura Penin., p. 101, pl. VI, figs. 16, 17. Moll. Rem. Uppern. Part, p. 15, pl. I, fig. 4.

A few small examples.

Fossil occurrence.—*Shimo-Ogawa Region*: Matsugane Mine. Shirado-Beds. Musashinos.

Living.—Northern and Central Japan.

18. *Dentalium* sp.

CM 22490

A fragment which looks like *Dentalium octogonum* Lam. (Yokoyama,

Foss. Miura Penin., p. 103, pl. VI, figs. 22—24), but having more ribs (more than 10).

CM 22491

Fossil occurrence.—*Minato Region*: Ushikubo.

CM 22492

### 19. *Panope generosa*, Gould.

*Panope generosa*. Yokoyama, Foss. Up., Musash., p. 121, pl. V, figs. 14, 15. Foss. Moll. Neog. Izumo, p. 4. Tert. Moll. Dainichi, p. 14. Moll. Rem. Upper. Part, p. 16, pl. VI, fig. 6

Rather rare and mostly in fragments.

✓

Fossil occurrence.—*Minato Region*: Ushikubo. *Tateishi Region*: Between *Tateishi* and *Shiogu*. Pliocene of Izumo and Dainichi. Shirado-Beds. Upper Musashino.

Living.—Northern Japan. West coast of America.

CM 22493

### 20. *Mactra sulcataria*, Deshayes.

*Mactra sulcataria*. Yokoyama, Foss. Up. Musashino, p. 126, pl. VII, fig. 6. Moll. Rem. Upper. Part, p. 17.

Several examples.

Fossil occurrence.—*Minato Region*: Isozaki. Shirado-Beds. Upper Musashino.

Living.—Northern, Central and Western Japan. North China. Possiet Bay near Vladivostok.

CM 22494

### 21. *Spisula grayana*, (Schrenck).

*Spisula grayana*. Yokoyama, Moll. Rem. Upper. Part, p. 17, pl. II, fig. 11.

Several large specimens.

Fossil occurrence.—*Minato Region*: Isozaki. Shirado-Beds. Upper Musashino.

Living.—Northern Japan. Behring Sea.

CM 22495

### 22. *Raeta elliptica*, Yokoyama.

*Raeta elliptica*. Yokoyama, Foss. Up. Musashino, p. 131, pl. VIII, fig. 7.

A single small valve.

Fossil occurrence.—*Minato-Region*: Near Kaimon Bridge. Upper Musashino.

CM 22496

### 23. *Tresus* sp.

A cast looking like *Tresus nuttali* Conr. (Foss. Up. Musash., p. 133, pl. VIII, fig. 8) fossil as well as living in Japan.

Fossil occurrence.—*Tateishi Region*: Between *Tateishi* and *Shiogu*.

24. *Tellina* sp,

CM 22497

A cast transversely elongated and looking like *Tellina alternata* Say var. *chibana* Yok. (Moll. Rem. Lowest Part, pl. II, fig. 20), though not well determinable. The rock in which it is contained is a shale.

Fossil occurrence.—*Shimo-Ogawa Region*: Shibahara.

25. *Macoma inquinata*, (Deshayes).

CM 22498

*Macoma inquinata*. Yokoyama, Moll. Rem. Lowest Part, p. 13. Foss. Miura Penin., p. 117, pl. VIII, figs. 1, 2.

A single valve.

Fossil occurrence.—*Minato Region*: Ushikubo. Iwaki-Beds. Pliocene of Kii. Lower and Upper Musashino.

Living.—Northern, Central and Western Japan. Sea of Okhotsk. West coast of North America.

26. *Dosinia troscheli*, Lischke.

CM 22499

*Dosinia troscheli*. Yokoyama, Foss. Miura Penin., p. 119, pl. VIII, figs. 5, 6. Moll. Rem. Upper. Part, p. 20.

Rather rare.

Fossil occurrence.—*Minato-Region*: Isozaki, *Shimo-Ogawa Region*: Ochiai and Shibahara. *Tateishi Region*: Between Tateishi and Shiogu. Shirado-Beds. Pliocene of Dainichi. Lower and Upper Musashino.

Living.—Central and Western Japan.

CM CM27093

CM CM27094 (R)

34

CM CM27095

27. *Dosinia angulosa*, Philippi.

Pl. II. Figs. 19, 20.

*Dosinia angulosa*. Philippi, Catalogue, p. 126. Philippi, Abbild., II, p. 229, pl. VI, fig. 1. Iwakawa, Catal., p. 297.

*Artemis penicillata*. Reeve, Conch. Icon., fig. 33.

A species close to the preceding, but distinguished by its more circular, consequently higher shape. The length and height in this species are nearly equal and the angle formed by the postero-dorsal and posterior borders is not so marked as in *Dosinia troscheli*. Moreover the pallial sinus is more rounded at end and finger-like in this species. The largest example is a left valve 55 millim. in length and height. Not rare in the Minato Region.

Fossil occurrence.—*Minato-Region*: Isozaki, Ushikubo and Hiraiso. *Shimo-Ogawa Region*: Shibahara.

Living.—Western Japan. Philippines and further south.

CM 22500

CM 22501

CM 22502-2-19

CM 22503

CM 22504-2-20

CM 22505

CM 22506

CM 22507

28. *Venus (Mercenaria) stimpsoni*, Gould.

*Venus (Mercenaria) stimpsoni*. Yokoyama, Foss. Up. Musash., p. 148, pl. XI, figs. 11, 12. Moll. Rem. Upper. Part, p. 21.

Several large specimens, though all are casts.

Fossil occurrence.—*Tateishi Region*: Between Tateishi and Shiogu. *Shimo-Ogawa Region*: Matsugane Mine. Shirado-Beds. Pliocene of Izumo. Upper Musashino.

Living.—Northern and Western Japan.

CM 22508

CM 22509

CM 22510-3-13

CM 22511

29. *Tapes variegatus*, Hanley.

Pl. III. Fig. 13.

*Tapes variegatus*. Yokoyama, Foss. Miura Penin., p. IX, figs. 4, 5. Foss. Up. Musash., p. 153.

Several specimens, one of which is 40 millim. high. The sculpture is somewhat different from that of the living ones by having the impressed radiating lines at more regular intervals and running more or less straight (not so wavy as in the latter).

Fossil occurrence.—*Minato Region*: Ushikubo, Kaimon Bridge, and Isozaki.

Living.—Central and Western Japan. Philippines.

CM 22512

CM 22513

CM 22514

CM 22515

CM 22516

CM 22517-3-10, 11

30. *Cardium shinjiense*, Yokoyama.

Pl. III, Figs. 10, 11.

*Cardium shinjiense*. Yokoyama, Molluscan Rem. Lowest. Part, p. 16, pl. III, figs. 13-15. Moll. Rem. Up. Part, p. 23.

A few examples.

Fossil occurrence.—*Minato Region*: Ushikubo, Isozaki. *Shimo-Ogawa Region*: Ochiai (Shibahara), Shibahara (in a fine yellowish tuffaceous shale) and Tatehita-zawa (in a soft grey sandy shale). Iwaki and Asagai-Beds. Shirado-Beds. Pliocene of Izumo.

CM 22518

CM 22519

31. *Diplodonta usta*, Gould.

*Diplodonta usta*. Yokoyama, Foss. Up. Musash., p. 159 pl. XIII, fig. 3. Foss. Miura Penin., p. 130, pl. IX, figs. 14-16.

Rare.

Fossil occurrence.—*Shimo-Ogawa Region*: Ochiai (Shibahara). *Minato Region*: Kaimon Bridge, Minato. Musashinos.

Living.—Northern and Central Japan.

CM 22520

CM 22521-3-12.

CM 22522

CM 22523

CM 22524

CM 22525

CM 22526

CM 22527

32. *Lucina (Phacoides) borealis*, Linné.

Pl. III. Figs. 1, 2.

*Lucina (Phacoides) borealis*. Yokoyama, Moll. Rem. Upper. Part, p. 24, pl. V. figs. 5-8.

Quite common at Shimo-Ogawa, though mostly in small forms.

Fossil occurrence.—*Shimo-Ogawa Region*: Shibahara, Ochiai, Tate-shita-zawa (shale) and Tanishita. *Minato Region*: Kaimon Bridge, Minato. *Tateishi Region*: Between Tateishi and Shiogu. Shirado-Beds. Musashinos. Pliocene of Kii. Miocene, Pliocene and Pleistocene of Europe.

Living.—Central Japan. Atlantic.

CM 22528

### 33. *Venericardia ferruginea*, (A. Adams).

*Venericardia ferruginea*. Yokoyama, Moll. Rem. Upper. Part, p. 24, pl. V, fig. 4.

*Venericardia sp.* Yokoyama, Moll. Rem. Lowest Part, p. 19, pl. III, figs. 8, 9.

Rare.

Fossil occurrence.—*Minato Region*: Ushikubo. Iwaki-Beds. Shirado-Beds. Musashinos.

Living.—Northern Japan.

CM 22529

### 34. *Cardita cumingiana*, Dunker.

*Cardita cumingiana*. Yokoyama, Foss. Miura Penin., p. 137, pl. X, fig. 16, pl. XI, fig. 1.

A strongly worn specimen.

Fossil occurrence.—*Minato-Region*: Hiraiso. Musashino.

Living.—Central and Western Japan.

### 35. *Lima goliath*, Sowerby.

CM 22530

*Lima goliath*. Yokoyama, Moll. Rem. Upper. Part, p. 26, pl. III, figs. 1, 4.

A single valve.

Fossil occurrence.—*Tateishi Region*: Between Tateishi and Shiogu, Shirado-Beds. Lower Musashino.

Living.—Central Japan. Patagonia.

### 36. *Pecten laetus*, Gould.

CM 22531

*Pecten laetus*. Yokoyama, Moll. Rem. Upper. Part, p. 26. Foss. Miura Penin., p. 152, pl. XIV, figs. 1, 2. Foss. up. Musashi., p. 180, pl. XIV, fig. 20.

Only two isolated valves.

Fossil occurrence.—*Minato Region*: Hiraiso. Shirado-Beds. Musashinos. Pleistocene of Awa.

Living.—Northern, Central and Western Japan.

### 37. *Pecten iwakianus*, Yokoyama.

CM 22532-3-12

Pl. III. Fig. 13. ✓

A fragment of a large valve which, however, shows such a peculiar sculpture as not to be confounded with any hitherto described.

The shell is nearly 80 millim. long, strongly compressed and furnished with more than twenty elevated, rounded, radiating ribs, narrower than the valleys in each of which there is a fine riblet. The ribs as well as the riblets are closely scaled.

The species comes closest to *Pecten crassicostatus* Sow. (Yokoyama, Foss. Miura Penin., pl. XII, fig. 7) living and fossil in Japan. This latter, however, has the ribs broader than the valleys and more coarsely and distantly scaled.

Fossil occurrence.—*Tateishi Region*: Between Tateishi and Shiogu.

CM 22533

38. *Arca* sp.

A young form looking like that of *Arca subcrenata* Lke. (Yokoyama, Foss. Up. Musash., p. 187, pl. PV, fig. 12).

Fossil occurrence.—*Minato Region*: Ushikubo.

CM 22534

CM 22535

39. *Pectunculus vestitus*, Dunker.

*Pectunculus vestitus*. Yokoyama, Foss. Miura Penin., p. 167, pl. XVIII, figs. 10, 11. Foss. Up. Musash., p. 189, pl. XVI, figs. 1–3. Moll. Rem. Lowest Part, p. 21. Moll. Rem. Upper. Part, p. 29.

Several large specimens.

Fossil occurrence.—*Minato-Region*: Ushikubo and Hiraiso. Iwaki-Beds. Shirado-Beds. Musashinos.

Living.—Central Japan.

CM 22536

40. *Parallelodon obliquatus*, Yokoyama.

*Parallelodon obliquatus*. Yokoyama, Foss. Miura Penin., p. 167, pl. XVII, figs. 10, 11. Moll. Rem. Lowest Part, p. 21. Moll. Rem. Upper. Part, p. 29.

Several small examples.

Fossil occurrence.—*Minato Region*: Hiraiso. Shirado-Beds. Musashinos. Pleistocene of Awa.

Living.—Northern and Central Japan.

(M 22537)

41. *Yoldia sagittaria*, Yokoyama.

A single ill-preserved valve of the species described before (p. 10).

Fossil occurrence.—*Tateishi Region*: Between Tateishi and Shiogu. Mizunoya—Beds.

42. *Nucula mirabilis*, Adams et Reeve.

CM 22538-3-6

Pl. III. Fig. 6.

CM 22539

*Nucula mirabilis*. Yokoyama, Foss. Miura Penin., p. 180, pl. XIX, fig. 9. Moll. Rem. Upper. Part, p. 30.

Quite frequent, some attaining a comparatively large size (over 30 millim. in length).

Fossil occurrence.—*Shimo-Ogawa Region*: Ochiai (Shibahara). Shirado-Beds. Musashino. Pliocene of Izumo.

---

Correction.

While this paper was under press, Dr. Tokunaga, after a more thorough examination, found the Tateishi layers as an equivalent of the Shirado-Beds in horizon, just as their fossils show.

---


## INDEX

	Page.		Page.
Arca sp . . . . .	20	Natica janthostoma Desh . . . . .	14
Bembix crumpii Pils . . . . .	15	Natica sp . . . . .	6
Bembix sp . . . . .	15	Nucula eximia Yok . . . . .	11
Calyptrea mammularis Brod . . . . .	13	Nucula insignis Ad . . . . .	20
Cardita cuningiana Dkr . . . . .	19	Nucula mirabilis Ad. et Rve . . . . .	21
Cardium shinjiense Yok . . . . .	18	Panope generosa Gld . . . . .	16
Cassis japonica Rve . . . . .	13	Parallelodon obliquatus Yok . . . . .	20
Chlorostoma argyrostoma Gm . . . . .	15	Pecten crassicostatus Lke . . . . .	20
Chlorostoma sp . . . . .	15	Pecten iwakianus Yok . . . . .	19
Chrysodomus despectus L . . . . .	12	Pecten laetus Gld . . . . .	19
Chrysodomus phoeniceus Dall . . . . .	12	Pecten tairanus Yok . . . . .	8
Crepidula aculeata Gm . . . . .	13	Pectunculus sp . . . . .	9
Crepidula convexa Yok . . . . .	14	Pectunculus vestitus DKr . . . . .	20
Crepidula ungana Dall . . . . .	14	Pleurotoma sp . . . . .	12
Dentalium sp . . . . .	15	Polinices ampla Rve . . . . .	14
Dentalium octogonum Lam . . . . .	15	Priene oregonensis Redf . . . . .	12
Diplodonta usta Ged . . . . .	18	Raeta elliptica Yok . . . . .	16
Dosinia angulosa Phil . . . . .	17	Ringicula sp . . . . .	11
Dosinia troscheli Lke . . . . .	17	Solemya tokunagai Yok . . . . .	11
Fissuridea humilis Yok . . . . .	15	Spisula grayana Schr . . . . .	16
Fissuridea patagonica Orb . . . . .	15	Tapes variegatus Hanl . . . . .	18
Fusus perplexus Rve . . . . .	12	Tellina alternata Say . . . . .	17
Helcioniscus pallidus Gld . . . . .	15	Tellina besshoensis Yok . . . . .	6
Leda inermis Yok . . . . .	9	Tellina sp . . . . .	6,17
Leda pennula Yok . . . . .	9	Tresus nuttali Conr . . . . .	16
Leda ramsayi Sm . . . . .	9	Tresus sp . . . . .	16
Lima goliath Sow . . . . .	19	Triton oregonensis Redf . . . . .	12
Lucina (Phacoides) borealis L . . . . .	7,18	Turritella nipponica Yok . . . . .	5,13
Lucina (Phacoides) sp . . . . .	7	Turritella sp . . . . .	6
Macoma dissimilis Mart . . . . .	6	Venericardia ferruginea Ad . . . . .	8,19
Macoma inquinata Desh . . . . .	17	Venericardia laxata Yok . . . . .	7
Mactra sulcataria Desh . . . . .	16	Venericardia orbica Yok . . . . .	8
Mercenaria stimpsoni Gld . . . . .	18	Venericardia pacifera Yok . . . . .	8
Nassa (Hima) japonica Ad . . . . .	12	Venus stimpsoni Gld . . . . .	18
		Yoldia sagittaria Yok . . . . .	10,20
		Yoldia tokunagai Yok . . . . .	10



Plate I.


- Fig. 1. *Turritella* sp. Mizunoya-Beds. Nagako, Kubota. P. 6.  
Fig. 2. *Lucina* (*Phacoides*) *borealis* L. (?). Kamenoo-Beds. Fukuda. P. 7.  
Figs. 3, 4, 5, 18. *Lucina* (*Phacoides*) sp. Kamenoo-Beds. 3. Ishimoriyama.  
4. Ōdate, Taira. 5. Ishibatake, Nishiki. 18. Tabasaka, Yumoto. P. 7.  
Figs. 6, 7. *Venericardia ferruginea* Ad. Kamenoo-Beds. Tsuchibashi,  
Kadono. P. 7.  
Figs. 8, 9. *Pecten tairanus* Yok. Somewhat enlarged. Kamenoo-Beds.  
8. Left valve. 9. Right valve. Kubo-Street, Taira. P. 8.  
Fig. 10. *Venericardia orbica* Yok. Kamenoo-Beds. Ishibatake, Nishiki. P. 8.  
Fig. 11. *Venericardia laxata* Yok. Kamenoo-Beds. Shimo-Yunagaya, Yumoto.  
P. 7.  
Fig. 12. *Venericardia laxata* Yok. (?) Kamenoo-Beds. Ebata, Yamada. P. 7.  
Fig. 13. *Venericardia pacifera* Yok. (?) Kamenoo-Beds. Daikatsu bore-hole,  
Yokonesawa, Uchigō. P. 8.  
Figs. 14, 15, 16. *Nucula eximia*, Yok. Kamenoo-Beds. 14. Kamenosawa,  
Mizunoya. 15. Hayashizaku, Iino. 16. Saku, Iino. P. 11.  
Fig. 17. *Pectunculus* sp. Kamenoo-Beds. Miyanosaku, Shimoyata. P. 9.  
Fig. 19. *Macoma dissimilis* Mart. (?) Kamenoo-Beds. Yunami. P. 6.  
Figs. 20, 21. *Solemya tokunagai* Yok. Kamenoo-Beds. Fukuda (west of).  
P. 11.



M. Yokoyama:—Molluscan Remains from the Jō-Ban Coal-Field.

## Plate II.


- Figs. 1—6. *Leda inermis* Yok. Kamenoo-Beds. 1, 2. Inouye, Uyeda. 3, 5. Osaku, Nakoso. 4. Tamatsuya, Izumi. 8. Hirakata Coast. P. 9.
- Figs. 7, 8, 9. *Leda pennula* Yok. Kamenoo-Beds. 7. Saku, Iino. P. 8. Takinosaku, Iino. 9. Daikatsu bore-hole, Yokonesawa. P. 9.
- Figs. 10, 11. *Yoldia sagittaria* Yok. Mizunoya-Beds. 10. Nagako, Kubota. 11. Between Kosai and Aburahira, Sekimoto. P. 10.
- Figs. 12—18. *Yoldia tokunagai* Yok. Kamenoo-Beds. 12. Honya, Izumi. 13. Takinosaku. 14. Ishimoriyama. 15, 18. Saku. 16. Miyanosaku. 17. Kami-Kabeya. P. 10.
- Figs. 19, 20. *Dosinia angulosa* Phil. Minato-Beds. 19. Hiraiso. 20. Shibahara. p. 17.



M. Yokoyama:—Molluscan Remains from the Jô-Ban Coal-Field.

### Plate III.

- Figs. 1, 2. *Lucina* (*Phacoides*) *borealis* L. Minato-Beds. Shibahara, Ochiai, Shimo-Ogawa. P. 18.
- Fig. 3. *Crepidula convexa* Yok. a. Dorsal view. b. Lateral view. Minato. P. 14.
- Figs. 4, 5. *Turritella nipponica* Yok. Minato-Beds. Ushikubo, Minato. P. 13.
- Fig. 6. *Nucula mirabilis* Ad. et Rve. Minato-Beds. Ochiai, Shimo-Ogawa. P. 21.
- Fig. 7. *Fissuridea humilis* Yok. a. b. Apical and lateral views. Minato-Beds. Hiraiso. P. 15.
- Figs. 8, 9. *Crepidula aculeata* Gm. 8. A worn specimen. 9. A somewhat better one. Minato-Beds. Hiraiso. P. 13.
- Figs. 10, 11. *Cardium shinjiense* Yok. Minato-Beds. Shibahara, Shimo-Ogawa. P. 18.
- Fig. 12. *Pecten iwakianus* Yok. Minato-Beds? Between Tateishi and Shiogu. P. 19.
- Fig. 13. *Tapes variegatus* Hant. Minato-Beds. Isozaki. P. 18.



M. Yokoyama:—Molluscan Remains from the Jô-Ban Coal-Field.

Vol. XLV., Art. 7, published June 25th,  
1925.

*Price in Tokyo, Yen 1.00.*

---

This Journal is on sale at  
MARUZEN CO., LTD., Tori Sanchoe, Nihonbashi, Tokyo.  
R. FRIEDLÄNDER u. SOHN, Carlstr. 11, Berlin, NW. 6.

---

大正十四年六月二十二日印刷  
大正十四年六月二十五日發行

編纂兼發行者 東京帝國大學

印刷者

東京市日本橋區兜町二番地  
星野錫

印刷所

東京市日本橋區兜町二番地  
東京印刷株式會社

賣捌所

東京市日本橋區通三丁目十四番地  
丸善株式會社



## NOTICE

### Vol. XLIII. (Concluded):

- Art. 1. K. YENDO:—A Monograph of the Genus *Alaria*. With 19 plates. Publ. May 30th, 1919.
- Art. 2. T. MATSUSHIMA:—Untersuchungen über die Wasseraufnahme bei abgeschnittenen Zweigen. Mit 2 Textfiguren. Publ. July 30th, 1919.
- Art. 3. Y. YOSHII:—Oekologische Studien über Vegetation der Ota Dünen. Mit 2 Tafeln und 8 Textfiguren. Publ. July 30th, 1919.
- Art. 4. Y. EMOTO:—Über die relative Wirksamkeit von Kreuz- und Selbstbefruchtung bei einigen Pflanzen. Mit 2 Tafeln und 6 Textfiguren. Publ. March 15th, 1920.
- Art. 5. T. KATŌ:—A Contribution to the Knowledge of the Cassiterite Veins of Pneumato-Hydrothermal or Hydrothermal Origin. A Study of the Copper-Tin Veins of the Abenobé District in the Province of Tajima, Japan. With 7 plates and 1 text figure. Publ. May 10th, 1920.
- Art. 6. S. TSUBOI:—Volcano Ōshima, Idzu. With 6 plates and 42 text figures. Publ. May 10th, 1920.
- Art. 7. M. TAHARA:—Cytologische Studien an einigen Kompositen. Mit 4 Tafeln und 15 Textfiguren. Publ. March 30th, 1921.
- Art. 8. Y. KUDŌ:—Enumeratio Labiatarum speciarum varietatum formarumque in Insulis Kurilensibus et Insula Yezoensi sponte nascentium. Publ. March 30th, 1921.

### Vol. XLIV. (Concluded):

- Art. 1. M. YOKOYAMA:—Fossils from the Upper Musashino of Kazusa and Shimosa. With 17 plates. Publ. July 7th, 1922.
- Art. 2. T. KABURAKI:—On some Japanese Freshwater Triclad, with a Note on the Parallelism in their Distribution in Europe and Japan. With 1 plate and 14 text figures. Publ. June 23rd, 1922.
- Art. 3. T. KABURAKI:—On some Japanese Tricladida Maricola, with a Note on the Classification of the Group. With 1 plate and 6 text figures. Publ. September 7th, 1922.
- Art. 4. T. KABURAKI:—On the Terrestrial Planarians from Japanese Territories. With 1 plate and 23 text figures. Publ. September 30th, 1922.
- Art. 5. T. TAKAGI:—Ueber das Reciprocitätsgesetz in einem beliebigen algebraischen Zahlkörper. Publ. September 16th, 1922.
- Art. 6. T. TERADA:—On some Remarkable Relations between the Yearly Variations of Terrestrial Phenomena and Solar Activities. With 17 text figures. Publ. January 24th, 1923.
- Art. 7. M. YOKOYAMA:—On some Fossil Shells from the Island of Saishū in the Strait of Tsushima. With 1 plate. Publ. January 24th, 1923.
- Art. 8. E. STECHOW:—Die Hydroidenfauna der Japanischen Region. Publ. February 26th, 1923.

### Vol. XLV.:

- Art. 1. M. YOKOYAMA:—Mollusca from the Coral-Bed of Awa. With 5 plates. Publ. October 3rd, 1924.
- Art. 2. M. YOKOYAMA:—Tertiary Mollusca from Dainichi in Tōtōmi. With 2 plates. Publ. March 31st, 1925.
- Art. 3. M. YOKOYAMA:—Molluscan Remains from the Lowest Part of the Jō-Ban Coal Field. With 5 plates. Publ. December 30th, 1924.
- Art. 4. Y. OZAWA:—On the Classification of Fusulinidae. With 4 plates and 3 text figures. Publ. February 22nd, 1925.
- Art. 5. M. YOKOYAMA:—Molluscan Remains from the Uppermost Part of the Jō-Ban Coal-Field. With 6 plates. Publ. March 21st, 1925.
- Art. 6. Y. OZAWA:—Paleontological and Stratigraphical Studies on the Permo-Carboniferous Limestone of Nāgato. Part II. Paleontology. With 14 plates. Publ. June 4th, 1925.
- Art. 7. M. YOKOYAMA:—Molluscan Remains from the Middle Part of the Jō-Ban Coal-Field. With 3 plates. Publ. June 25th, 1925.